

©2000

HOMOTOPY RINEHART COHOMOLOGY OF HOMOTOPY LIE-RINEHART PAIRS

LARS KJESETH

*(communicated by Larry Lambe)**Abstract*

We define homotopy Lie-Rinehart pairs and the associated homotopy Rinehart algebra in the context of coalgebras. We define homotopy Lie-Rinehart resolutions and present conditions under which the associated homotopy Rinehart algebra is a cohomological model for the Rinehart algebra of the resolved Lie-Rinehart pair.

1. Introduction

The Rinehart cohomology of homotopy Lie-Rinehart resolutions of Lie-Rinehart pairs arose from the study of the BFV (Batalin, Fradkin and Vilkovisky, [FV75], [BV77], [BF83], [BV83] and [BV85]) formulation of classical BRST cohomology (Becchi, Rouet and Stora [BRS75] and, independently, Tyutin [Tyu75]). The classical BRST algebra $(\mathcal{A}, \mathcal{D})$ (see, for example, [Kim92b], [Kim93], [Sta92], [Kim92a], [FHST89], [Sta88], [Sta96], [KS87] and [HT92]) is a differential graded Poisson algebra which, in certain cases, is a *cohomological model* for the Rinehart algebra $(Alt_{\mathbf{B}}(\mathfrak{g}, \mathbf{B}), \delta_R)$ of alternating \mathbf{B} -multilinear functions from a Lie algebra \mathfrak{g} into the associative algebra \mathbf{B} [Rin63]. The Rinehart complex is a subcomplex of the Chevalley-Eilenberg complex ([CE48]) precisely when $(\mathbf{B}, \mathfrak{g})$ is a *Lie-Rinehart pair* satisfying certain *Lie-Rinehart relations* (see below).

In homological algebra, a traditional strategy is to replace both \mathbf{B} and \mathfrak{g} in $Alt_{\mathbf{B}}(\mathfrak{g}, \mathbf{B})$ with free or projective resolutions $\mathbf{R}_{\mathbf{B}}$ and $\mathbf{R}_{\mathfrak{g}}$ which have the same or similar algebraic structures, at least up to homotopy. The new object $Alt_{\mathbf{R}_{\mathbf{B}}}(\mathbf{R}_{\mathfrak{g}}, \mathbf{R}_{\mathbf{B}})$ is a cohomological model for $Alt_{\mathbf{B}}(\mathfrak{g}, \mathbf{B})$ if $Alt_{\mathbf{R}_{\mathbf{B}}}(\mathbf{R}_{\mathfrak{g}}, \mathbf{R}_{\mathbf{B}})$ is a differential graded commutative algebra (a dcca) with the same cohomology as $Alt_{\mathbf{B}}(\mathfrak{g}, \mathbf{B})$. In contrast, the BFV construction of the classical BRST algebra (see, for example, [Kim93]) begins by replacing \mathbf{B} with the Koszul-Tate resolution [Tat57], but side-steps replacing \mathfrak{g} with a suitable resolution, opting instead to adjoin formal (ghost) variables to the Koszul-Tate resolution. They then exploit a graded Poisson bracket to construct a differential. Attempting to construct the dcca $Alt_{\mathbf{R}_{\mathbf{B}}}(\mathbf{R}_{\mathfrak{g}}, \mathbf{R}_{\mathbf{B}})$ fails if we resolve \mathbf{B} and \mathfrak{g} without preserving (as far as possible) the Lie-Rinehart structure of the pair $(\mathbf{B}, \mathfrak{g})$. Our view in this paper is that a model for a Rinehart complex should be Rinehart-like. But to resolve a Lie-Rinehart pair with a new pair $(\mathbf{R}_{\mathbf{B}}, \mathbf{R}_{\mathfrak{g}})$ which is Lie-Rinehart, we must give up a strictly Lie structure on $\mathbf{R}_{\mathfrak{g}}$ in favor of a strongly homotopy Lie structure.

In its earliest incarnation, [Sul77] and [SS], a strongly homotopy Lie (shLie) algebras was defined on the tensor coalgebra $T^c L$ of a graded module L . In [LM95] and [LS], the shLie structure was shifted (almost by brute force) onto the graded symmetric coalgebra on the suspension of L , that is, on $\bigwedge(sL)$. Parts of this transition were quite rough and we felt there must be smoother means and more satisfying motivations for making the move. By taking a close look at choices of grading and specific actions of the symmetric group on a graded vector

Received 8 January 2001, revised ????; published on ????

2000 Mathematics Subject Classification: primary 17B55, 17B81; secondary 17B56, 17B99

Key words and phrases: strongly homotopy Lie algebras, Lie-Rinehart pairs, Rinehart cohomology, BRST cohomology

© 2001, Lars Kjeseth. Permission to copy for private use granted.

space, we discovered we could identify a subcoalgebra $\bigwedge L$ of $T^c L$, which is isomorphic to $\bigwedge(sL)$ and on which an shLie structure can be defined as it was on $T^c L$ in [LM95] and [LS] (See §2).

Grading and Sign Conventions

In order to understand the language of this paper, we must establish our grading and sign conventions first.

All vector spaces, algebras and coalgebras in this paper are over a field k of characteristic zero. All tensor products are over k . All maps are at least k -linear or k -multilinear.

Let $V = \{V_p\}_0^\infty$ be a graded vector space over k or, more generally, a graded module over an algebra A over k . There are four distinct gradings on the tensor module $\bigotimes V = \{V^{\otimes n}\}_{n=0}^\infty$: the *external*, *internal*, *combined* and *total* gradings. We shall denote $v_1 \cdots v_n \in V^{\otimes n}$ by $v_{[1 \text{ to } n]}$. The *external degree*, $\mathbf{ed}(v_{[1 \text{ to } n]})$ is given by the number n of tensor components. The *internal degree*, $\mathbf{id}(v_{[1 \text{ to } n]})$ comes from the internal grading on V ; each tensor component v_i is an element of V_{p_i} for some p_i . The internal degree is the sum $\sum_{i=1}^n p_i$. The *combined degree* $\|v_{[1 \text{ to } n]}\|$ is the vector $\langle \mathbf{ed}(v_{[1 \text{ to } n]}), \mathbf{id}(v_{[1 \text{ to } n]}) \rangle$. Finally, for any bigraded object, the *total degree* $\mathbf{td}(v_{[1 \text{ to } n]})$ is the sum $\mathbf{ed}(v_{[1 \text{ to } n]}) + \mathbf{id}(v_{[1 \text{ to } n]})$.

A function $f : \bigotimes V \rightarrow \bigotimes V$ is of *homogeneous degree* r with respect to a specific grading if $f((\bigotimes V)_p) \subset (\bigotimes V)_{p+r}$ for all p . For each of the four gradings, the degree of the homogeneous map f is the difference between the degree of $v_{[1 \text{ to } n]} \in \bigotimes V$ and the degree of its non-zero image under f .

The Koszul sign convention states that exchanging two objects of homogeneous degrees p and q (whether elements or maps) introduces a factor of $(-1)^{pq}$. In this paper, we apply the Koszul sign convention consistently, no matter which grading. (In the case of the combined grading, the product of p and q will be the dot product of the degree vectors.)

Resolving a Lie algebra with an shLie algebra resolution is much simpler if we view the Lie algebra as a graded object. After all, since the shLie algebra resolution is defined on a graded coalgebra, it makes sense that the original Lie algebra should be realized on a graded coalgebra as well, where, at the very least, every element has an external degree. We must nail down the relation between the symmetry of the Lie bracket and the degree we assign to the elements of the Lie algebra. If each element of the Lie algebra \mathfrak{g} is assigned an external degree of one and an internal degree of zero, we have essentially three choices for describing \mathfrak{g} as a coalgebra. In each case, the skew-commutativity of l_2 appears as the result of applying the Koszul sign convention. If we select the natural bigrading on $T^c \mathfrak{g}$, the element x has bidegree $\langle 1, 0 \rangle$, which we treat as a vector, and

$$l_2(x \ y) = (-1)^{\langle 1,0 \rangle \cdot \langle 1,0 \rangle} l_2(y \ x) = -l_2(y \ x).$$

We might instead choose the total degree, in which case the element x has degree $0 + 1 = 1$. Here again,

$$l_2(x \ y) = (-1)^{1 \cdot 1} l_2(y \ x) = -l_2(y \ x).$$

Finally, if we consider the tensor coalgebra on the suspension of \mathfrak{g} under the internal grading, then the element x again has degree 1 and l_2 again appears skew-commutative, but is actually graded commutative.

When it resolves a Lie algebra, an shLie algebra must operate as a differential object. The differential must have degree -1 with respect to a single grading. So the shLie algebra resolution $\mathbf{R}_{\mathfrak{g}}$ of \mathfrak{g} must be graded either with respect to the total degree or with respect to the suspended internal degree. Since the differential graded commutative (dgc) algebra $\mathbf{R}_{\mathbf{B}}$ which resolves \mathbf{B} is graded by internal degree, the suspended internal grading for the shLie algebra resolution is the preferred choice. Therefore, we will also view the Lie algebra \mathfrak{g} in its suspended form \mathfrak{sg} .

Background

For any smooth manifold M , the set of smooth functions $C^\infty(M)$ is an associative commutative algebra and the set of smooth vector fields on the manifold, $\Gamma(TM)$, forms a Lie algebra. Both $C^\infty(M)$ and $\Gamma(TM)$ are modules over each other. Moreover, the the module actions satisfy the equations $(fX) \cdot g = f(X \cdot g)$ and $[X, fY] = f[X, Y] + (X \cdot f)Y$ for all $f, g \in C^\infty(M)$ and $X, Y \in \Gamma(TM)$. A Lie-Rinehart pair is a couple $(\mathbf{B}, \mathfrak{g})$ which admits the analogous structure, where \mathfrak{g} is a Lie algebra (we will suspend it later), \mathbf{B} is an algebra, and both are modules over each other.

Definition 1.1. : [Rin63] We denote the left \mathbf{B} -module action μ on \mathfrak{g} by $\mu(a \ \alpha) := a\alpha$. Let $\omega : \mathfrak{g} \ \mathbf{B} \rightarrow \mathbf{B}$ (or, alternatively, $\omega : \mathfrak{g} \rightarrow \text{Der}(\mathbf{B})$) denote the \mathfrak{g} -module action on \mathbf{B} . The pair $(\mathbf{B}, \mathfrak{g})$ is a Lie-Rinehart pair, provided the Lie-Rinehart relations (LRa) and (LRb) are satisfied for all $a, b \in \mathbf{B}$ and $x, y \in \mathfrak{g}$:

- LRa: $\omega(ax \ b) = a \cdot \omega(x \ b)$, where \cdot indicates the multiplication on \mathbf{B} .
- LRb: $[x, ay] = a[x, y] + \omega(x \ a)y$.

The term Lie-Rinehart pair is not widely used. More often, \mathfrak{g} has been called a (\mathbf{B}, k) -Lie algebra, [Rin63] [Pal61] and [Her72]. More recently, Lie-Rinehart pairs have appeared as Lie algebroids (see [dSW]).

The first Lie-Rinehart relation (LRa) states that ω is \mathbf{B} -linear in \mathfrak{g} .

Suppose \mathfrak{g} is finitely generated over \mathbf{B} , that is to say, with respect to a generating set $\{X_\alpha\}$, every element x of \mathfrak{g} can be expressed as the sum $b^\alpha X_\alpha$ where $b^\alpha \in \mathbf{B}$. (We use the Einstein summation convention throughout.) The second Lie-Rinehart relation (LRb) and its skew-symmetric counterpart, $[ax, y] = a[x, y] - \omega(y \ a)x$, imply that for any element $b^\alpha X_\alpha \ b^\beta X_\beta$ of $\mathfrak{g} \ \mathfrak{g}$ the bracket must have the form

$$[b^\alpha X_\alpha, b^\beta X_\beta] = b^\alpha b^\beta [X_\alpha, X_\beta] + \omega(b^\alpha X_\alpha \ b^\beta)X_\beta - \omega(b^\beta X_\beta \ b^\alpha)X_\alpha. \tag{1}$$

The second Lie-Rinehart relation (LRb) is equivalent to equation (1).

For any Lie algebra \mathfrak{g} and \mathfrak{g} -module \mathbf{B} , an n -multilinear function $f_n : \mathfrak{g}^{\times n} \rightarrow \mathbf{B}$ is alternating if

$$f_n(x_1, \dots, x_i, x_{i+1}, \dots, x_n) = -f_n(x_1, \dots, x_{i+1}, x_i, \dots, x_n).$$

The Chevalley-Eilenberg complex is the set of all alternating multilinear functions $Alt_k(\mathfrak{g}, \mathbf{B})$, graded by n , and equipped with a degree +1 differential $\delta_{CE} : Alt_k^n(\mathfrak{g}, \mathbf{B}) \rightarrow Alt_k^{n+1}(\mathfrak{g}, \mathbf{B})$ given by

$$\begin{aligned} \delta_{CE} f_n(x_0, \dots, x_n) &= \sum_{i=0}^n (-1)^i \omega(x_i \ f_n(x_0, \dots, \widehat{x}_i, \dots, x_n)) \\ &\quad - \sum_{i < j} (-1)^{i+j-1} f_n([x_i, x_j], x_0, \dots, \widehat{x}_i, \dots, \widehat{x}_j, \dots, x_{\sigma(n+1)}), \end{aligned} \tag{2}$$

where \widehat{x}_k indicates that x_k should be omitted. Any element $b \in \mathbf{B}$ is considered a 0-cochain. The image of b under δ_{CE} is defined by setting $\delta_{CE} b(x) = \omega(x \ b)$. When \mathbf{B} is an algebra, the Chevalley-Eilenberg complex is a differential graded commutative algebra. For f_n and g_m in $Alt_k(\mathfrak{g}, \mathbf{B})$, the product $f_n \smile g_m$ is given by

$$(f_n \smile g_m)(x_1, \dots, x_{n+m}) = \sum_{\substack{\sigma \\ (n,m) \\ \text{unshuffles}}} f_n(x_{\sigma(1)}, \dots, x_{\sigma(n)}) g_m(x_{\sigma(n+1)}, \dots, x_{\sigma(n+m)}).$$

An (n, m) -unshuffle is any permutation σ in the symmetric group Σ_{n+m} such that

$$\underbrace{\sigma(1) < \dots < \sigma(n)}_{\text{first } \sigma \text{ hand}} \quad \text{and} \quad \underbrace{\sigma(n+1) < \dots < \sigma(n+m)}_{\text{second } \sigma \text{ hand}},$$

where $\sigma(j)$ is the element of the set $\{1, \dots, n + m\}$ moved to the j^{th} position under σ . The differential δ_{CE} acts as a derivation with respect to this multiplication. The cohomology of this complex with respect to δ_{CE} is the Chevalley-Eilenberg cohomology of \mathfrak{g} with coefficients in \mathbf{B} [CE48].

Suppose $(\mathbf{B}, \mathfrak{g})$ is a Lie-Rinehart pair and the alternating function f_n is \mathbf{B} -multilinear, i.e., $f_n(a_1x_1, \dots, a_nx_n) = a_1 \cdots a_n f_n(x_1, \dots, x_n)$. Because the Lie action map ω maps \mathfrak{g} into the derivations of \mathbf{B} and as a result of the Lie-Rinehart relations, the image of f_n under the Chevalley-Eilenberg differential δ_{CE} is again \mathbf{B} -multilinear, despite the fact that the bracket is not \mathbf{B} -multilinear. The Rinehart algebra $R = \text{Alt}_{\mathbf{B}}(\mathfrak{g}, \mathbf{B})$ with differential $\delta_R = \delta_{CE}$ is a subcomplex of the Chevalley-Eilenberg algebra and the cohomology with respect to δ_R is the Rinehart cohomology of \mathfrak{g} with coefficients in \mathbf{B} [Rin63].

Summary

First, we define the Lie algebras of subordinate derivation sources, resting coderivations and shared Lie modules (§2). Following the lead of [LM95] and [LS], we define both homotopy and non-homotopy Lie algebras, Lie algebra modules, Lie-Rinehart pairs and Rinehart cohomology, all in the coalgebra setting (§§3 and 4). We piece together homotopy Lie-Rinehart resolutions for Lie-Rinehart pairs and present conditions under which the homotopy Rinehart algebra for a homotopy Lie-Rinehart resolution is a model for the Rinehart cohomology complex for a Lie-Rinehart pair (§5).

We have omitted all sign arguments from the proofs in this paper, as they are not generally instructive.

2. Subordinate derivation sources, resting coderivations and shared Lie modules

Motivation

The following brief and not unduly precise reexamination of the Lie-Rinehart relations and the Rinehart differential provides motivation for why coalgebras are an appropriate setting for both the homotopy and the non-homotopy versions of Lie algebras, Lie modules, Chevalley-Eilenberg cohomology and Rinehart cohomology. In particular, the discussion below highlights why we define *subordinate* and *resting* coderivations, as well as *shared* Lie modules.

The underlying module structure for the tensor coalgebra $T^c \mathfrak{g}$ on \mathfrak{g} is $\bigotimes \mathfrak{g}$, which is the collection $\{\mathfrak{g}^{\otimes n}\}_{n=0}^{\infty}$.

We rename the bracket as l_2 and extend l_2 to a map from $\mathfrak{g}^{\otimes n} \rightarrow \mathfrak{g}^{\otimes n-1}$ for any $n > 2$ by setting (up to sign)

$$l_2(x_1 \otimes \cdots \otimes x_n) = \sum_{\sigma \in (2, n-2)} \pm l_2(x_{\sigma(1)} \otimes x_{\sigma(2)} \otimes x_{\sigma(3)} \otimes \cdots \otimes x_{\sigma(n)}), \tag{3}$$

where the sum runs over all $(2, n - 2)$ -unshuffles σ . The bracket l_2 is now a *coderivation* on $T^c \mathfrak{g}$, although it has lost its skew-commutativity. We regain the symmetry by passing to an appropriate subcoalgebra. In this context, we express the Jacobi identity by $l_2 \circ l_2 = 0$. Or, since the set of coderivations on a coalgebra forms a Lie algebra, the Jacobi identity is equivalent to $[l_2, l_2] = 0$.

Similarly, we rename the Lie action map ω as m_2 and extend m_2 as a coderivation on $T^c \mathfrak{g} \otimes \mathbf{B}$ which is *subordinate* to l_2 by setting

$$m_2(x_1 \otimes \cdots \otimes x_n \otimes b) = l_2(x_1 \otimes \cdots \otimes x_n) \otimes b \pm \sum_{\rho \in (n-1, 1)} \pm x_{\rho(1)} \otimes \cdots \otimes x_{\rho(n-1)} \otimes m_2(x_{\rho(n)} \otimes b) \tag{4}$$

for any $n > 1$, where l_2 has been extended as in equation (3). Passing again to the appropriate subcoalgebra, we regain the desired symmetry. The condition any Lie action must satisfy becomes $m_2 \circ m_2 = 0$ or $[m_2, m_2] = 0$.

For the Lie-Rinehart pair $(\mathbf{B}, \mathfrak{g})$, the Lie algebra is a module over \mathbf{B} , but the bracket l_2 is not \mathbf{B} -linear. Rather, the second Lie-Rinehart relation states that $l_2(ax \quad by)$ consists of a \mathbf{B} -linear piece

$$abl_2(x \quad y),$$

and two other terms

$$m_2(ax \quad b)y - m_2(by \quad a)x$$

(compare with equation 1). Any coderivation which shares this property with a specific subordinate coderivation is called a *resting* coderivation, e.g., l_2 rests on m_2 .

Finally, we extend an alternating function $f_n : \mathfrak{g}^n \rightarrow \mathbf{B}$ to a new alternating function f_n from $T^c \mathfrak{g}$ into $T^c \mathfrak{g} \otimes \mathbf{B}$ by setting $f_n(x_1 \quad \dots \quad x_{n+k}) =$

$$\sum_{\substack{\sigma \\ (k,n)}} \pm x_{\sigma(1)} \quad \dots \quad x_{\sigma(k)} \quad f_n(x_{\sigma(k+1)} \quad \dots \quad x_{\sigma(k+n)}) \tag{5}$$

on \mathfrak{g}^{k+n} and then passing to the appropriate subcoalgebra. Using equations (3), (4) and (5), the Chevalley-Eilenberg differential (and hence, the Rinehart differential) becomes

$$\delta_{CE} f_n = m_2 \circ f_n \pm f_n \circ l_2,$$

which looks remarkably like some sort of commutator bracket action.

In our more rigorous discussion of these ideas, we focus on the following algebraic objects.

1. The graded Lie algebra $\text{Coder}_W^W(\wedge(sV))$ is the set of all coderivations on the free graded commutative algebra $\wedge(sV)$, each of which rests on its own specific subordinate coderivation in the Lie algebra $\text{Coder}_W^W(\wedge(sV) \quad W)$ (see 2).
2. For a graded commutative algebra W , the Lie algebra $\text{Coder}_W^W(\wedge(sV) \quad W)$ is the set of all coderivations, each of which is subordinate to a specific resting coderivation and is also a W -derivation source. A W -derivation source m is the extension as a coderivation onto $\wedge(sV)$ of a map m on $(sV)^{\wedge n} \quad W$ which acts like a Lie module structure map, i.e., $m : (sV)^{\wedge n} \rightarrow \text{Der}(W)$.
3. The graded commutative algebra $\text{Hom}_W(\wedge(sV), \wedge(sV) \quad W)$ is the set of all W -linear maps on $\wedge(sV)$ with coefficients in $\wedge(sV) \quad W$ and it admits a shared Lie module structure over both $\text{Coder}_W^W(\wedge(sV) \quad W)$ and $\text{Coder}_W^W(\wedge(sV))$.

We begin our more formal treatment with a return to the basics.

Coalgebras and Subcoalgebras

A *graded coassociative coalgebra* is a pair (\mathbf{C}, Δ) , where \mathbf{C} is a graded module over k together with a 0-degree coassociative comultiplication $\Delta : \mathbf{C} \rightarrow \mathbf{C} \otimes \mathbf{C}$. The comultiplication breaks up into pieces $\Delta_{p,q} : \mathbf{C}^{p+q} \rightarrow \mathbf{C}^p \otimes \mathbf{C}^q$, based on the bigrading of the range. A function f with degree $|f| = r$ is a *coderivation* on \mathbf{C} if, for all p and q , the diagram

$$\begin{array}{ccc} \mathbf{C}^{p+q} & \mathbf{C}^q \otimes \mathbf{C}^p & \mathbf{C}^{q+r} \xrightarrow{f \quad 1+1} f \quad \mathbf{C}^p \otimes \mathbf{C}^q \\ & \uparrow \Delta_{p+r,q} + \Delta_{p,q+r} & \uparrow \Delta_{p,q} \\ & \mathbf{C}^{p+q+r} & \xrightarrow{f} \mathbf{C}^{p+q} \end{array}$$

commutes, where again $(f \quad 1)(c_p \quad c_q) = f(c_p) \quad c_q$ and $(1 \quad f)(c_p \quad c_q) = (-1)^{r_p} c_p \quad f(c_q)$.

The set of all coderivations on a coalgebra C , denoted $\text{Coder}(C)$, is a graded Lie algebra under the graded commutator bracket, that is to say, $[f, g] = fg - (-1)^{|f||g|}gf$ for all f and $g \in \text{Coder}(C)$, where $|f|$ and $|g|$ are the degrees of f and g . For a graded module V , the

tensor module $\otimes V = \{V^n\}$ is a graded coassociative coalgebra T^cV with the standard comultiplication $\Delta : T^cV \rightarrow T^cV \otimes T^cV$ given by

$$\Delta(v_{[1 \text{ to } n]}) = \sum_{j=0}^n (v_{[1 \text{ to } j]}) \otimes (v_{[j+1 \text{ to } n]}),$$

where $V^0 \approx k$ and the terms with $j = 0$ and $j = n$ are $1 \otimes v_{[1 \text{ to } n]}$ and $v_{[1 \text{ to } n]} \otimes 1$ respectively.

Actions of the symmetric group Σ_∞ on T^cV

An action ρ_n of the symmetric group Σ_n on V^n is any homomorphism

$$\rho_n : \Sigma_n \rightarrow \text{End}(V^n).$$

An action $\rho = \{\rho_n\}_{n=1}^\infty$ of $\{\Sigma_n\}_{n=1}^\infty$ on T^cV is defined in the obvious way. The ρ -invariant subspace of T^cV forms a subcoalgebra. The two actions and associated subcoalgebras which play a role in this paper are listed below. The first one is the familiar graded commutative coalgebra $\wedge V$. The other one helps place this paper in its historical context.

1. The *internal graded* action ρ^\wedge recognizes only the internal grading on V^n and is given by $\sigma \cdot (v_{[1 \text{ to } n]}) = \text{Kid}(\sigma)v_{\sigma[1 \text{ to } n]}$ for all n , $\sigma \in \Sigma_n$ and $v_{[1 \text{ to } n]} \in V^n$. The symbol $v_{\sigma[1 \text{ to } n]}$ is shorthand for $v_{\sigma(1)} \cdots v_{\sigma(n)}$, where $\sigma(i)$ is the element of the ordered set $\{1, \dots, k\}$ which moves to the i^{th} position under σ . Since Σ_n acts transitively on itself, the ρ^\wedge -invariant subspace $\wedge V$ in T^cV is the *internal graded symmetric coalgebra* on V and is generated by elements of the form

$$\sum_{\sigma \in \Sigma_n} \text{Kid}(\sigma)v_{\sigma[1 \text{ to } n]},$$

which we will denote by $v_{[1 \text{ to } n]}^\wedge$. For all $v_{[1 \text{ to } n]}^\wedge \in V^{\wedge n}$ and $\sigma \in \Sigma_n$, we have the commutation relation $v_{[1 \text{ to } n]}^\wedge = \text{Kid}(\sigma)v_{\sigma[1 \text{ to } n]}^\wedge$. The coassociative comultiplication on $\wedge V$ is given by

$$\Delta(v_{[1 \text{ to } n]}^\wedge) = \sum_{j=0}^n \sum_{(j, n-j)\text{-unshuffle}} \text{Kid}(\rho) \left(v_{\rho[1 \text{ to } j]}^\wedge \right) \otimes \left(v_{\rho[j+1 \text{ to } n]}^\wedge \right).$$

2. The *combined graded* action ρ^\otimes is given by $\sigma \cdot (v_{[1 \text{ to } n]}) = \text{K}_{|| \ ||}(\sigma)v_{\sigma[1 \text{ to } n]}$. The ρ^\otimes -invariant subcoalgebra is the *combined graded symmetric coalgebra* $\bigotimes V$, generated by elements of the form

$$\sum_{\sigma \in \Sigma_n} \text{K}_{|| \ ||}(\sigma)v_{\sigma[1 \text{ to } n]}.$$

The commutation relation is $v_{[1 \text{ to } n]}^\otimes = \text{K}_{|| \ ||}(\sigma)v_{\sigma[1 \text{ to } n]}^\otimes$, and the comultiplication is given by

$$\Delta(v_{[1 \text{ to } n]}^\otimes) = \sum_{j=0}^n \sum_{(j, n-j)} \text{K}_{|| \ ||}(\rho) \left(v_{\rho[1 \text{ to } j]}^\otimes \right) \otimes \left(v_{\rho[j+1 \text{ to } n]}^\otimes \right),$$

where it is understood that the second sum is over all $(j, n - j)$ -unshuffles.

Let sV be the suspension of V , that is to say, $(sV)_p = V_{p-1}$ for all p . The subcoalgebra $\bigotimes V$ of the tensor coalgebra T^cV is isomorphic to the subcoalgebra $\wedge(sV)$ of $T^c(sV)$. This isomorphism replaces the bigrading on $\bigotimes V$ with its total grading, allowing it to function more easily as a differential object. The coalgebra $\wedge(sV)$ is graded by internal degree, which will

be the preferred setting when looking at graded maps from $\bigwedge(sV)$ into a differential graded commutative algebra, which are traditionally graded by internal degree.

The coalgebra isomorphism $S : \bigoplus V \rightarrow \bigwedge(sV)$ is not of homogeneous degree; each map $S^n : V^{\otimes n} \rightarrow (sV)^{\wedge n}$ increases the internal degree of an element by n . Exchanging a map f with S^n will produce a sign of $(-1)^{(0,n) \cdot \text{ed}(f), \text{id}(f)} = (-1)^{n \cdot \text{id}(f)}$. The inverse isomorphism of S^n shall be denoted by S^{-n} . The map S respects the appropriate actions of Σ_∞ .

Although we define action invariant maps on $T^c(sV)$, the definition applies to T^cV as well. Let W be a graded module.

Definition 2.1. : A map of graded modules $\widehat{l}_n : (sV)^{\wedge n} \rightarrow W$ is ρ_n -invariant if, for all $sv_{[1 \text{ to } n]}$ and $\sigma \in \Sigma_n$, $\widehat{l}_n(\sigma \cdot (sv_{[1 \text{ to } n]})) = \widehat{l}_n(sv_{[1 \text{ to } n]})$. Similarly, a map $\widehat{l} : T^c(sV) \rightarrow W$ is ρ -invariant if $\widehat{l}(\sigma \cdot (sv_{[1 \text{ to } n]})) = \widehat{l}(sv_{[1 \text{ to } n]})$ for all n and $\sigma \in \Sigma_n$.

If $\widehat{l} : T^c(sV) \rightarrow W$ is ρ -invariant, then \widehat{l} restricted to the ρ -invariant subcoalgebra is well-defined because Σ_n acts transitively on itself. The maps of greatest interest to us in this paper are either suspended internal graded symmetric (ρ^\wedge -invariant) or combined graded symmetric (ρ^\otimes -invariant). The map \widehat{l}_n is suspended internal graded symmetric if $\widehat{\text{Kid}}(\sigma)\widehat{l}_n(sv_{\sigma[1 \text{ to } n]}) = \widehat{l}_n(sv_{[1 \text{ to } n]})$ and l_n is combined graded symmetric if $\text{K}_{|| \ ||}(\sigma)l_n(v_{\sigma[1 \text{ to } n]}) = l_n(v_{[1 \text{ to } n]})$ for all $v_{[1 \text{ to } n]}$ and $\sigma \in \Sigma_n$. We do not lose any information if we restrict the map \widehat{l}_n to $(sV)^{\wedge n}$ (or restrict l_n to $V^{\otimes n}$) because

$$\begin{aligned} \widehat{l}_n(sv_{[1 \text{ to } n]}^\wedge) &= \widehat{l}_n\left(\sum_{\sigma \in \Sigma_n} \widehat{\text{Kid}}(\sigma)sv_{\sigma[1 \text{ to } n]}\right) \\ &= \sum_{\sigma \in \Sigma_n} \widehat{\text{Kid}}(\sigma)\widehat{l}_n(sv_{\sigma[1 \text{ to } n]}) \\ &= \sum_{\sigma \in \Sigma_n} \widehat{l}_n(sv_{[1 \text{ to } n]}) \\ &= n!\widehat{l}_n(sv_{[1 \text{ to } n]}). \end{aligned}$$

So whenever we consider collections of maps on T^cV with the same invariance, we pass to that invariant subcoalgebra. The isomorphism S implies that any map $l_n : V^{\otimes n} \rightarrow V$ induces a map $\widehat{l}_n : (sV)^{\wedge n} \rightarrow (sV)$ by setting $\widehat{l}_n = (-1)^{n \cdot \text{id}(l_n)}S^1l_nS^{-n}$. This process is invertible, so the set of all maps $l_n : V^{\otimes n} \rightarrow V$ is isomorphic to the set of all maps $\widehat{l}_n : (sV)^{\wedge n} \rightarrow sV$. Therefore, whatever we define and prove in the suspended internal graded setting has its analog in the combined graded setting, e.g., all Lie algebras and graded commutative algebra defined in the remainder of this section.

The graded Lie algebra $\text{Coder}(\bigwedge(sV))$

Following [Lad], one can extend any map $\widehat{l}_n : (sV)^{\wedge n} \rightarrow (sV)$ to a coderivation $\widehat{l}_n : T^c(sV) \rightarrow T^c(sV)$ by setting $\widehat{l}_n = 0$ on $(sV)^{\wedge k}$ for $k < n$ and, for $k \geq n$, setting

$$\widehat{l}_n(sv_{[1 \text{ to } k]}) = \sum_{i=1}^{k-n+1} (1 \quad i-1 \quad \widehat{l}_n \quad 1 \quad k-n+1-i)(sv_{[1 \text{ to } k]}).$$

Here, $(1 \quad i-1 \quad \widehat{l}_n \quad 1 \quad k-n+1-i)(v_{[1 \text{ to } k]}) =$

$$\widehat{\text{Kid}}(\widehat{l}_n; sv_{[1 \text{ to } i-1]})sv_{[1 \text{ to } i-1]} \quad \widehat{l}_n(sv_{[i \text{ to } i+n-1]}) \quad sv_{[i+n \text{ to } k]},$$

where $\widehat{\text{Kid}}(\widehat{l}_n; sv_{[1 \text{ to } i-1]})$ is the sign introduced by exchanging \widehat{l}_n with $sv_{[1 \text{ to } i-1]}$. However, extending \widehat{l}_n as a coderivation on $T^c(sV)$ does not guarantee that its invariance is preserved.

Exploiting the commutation relations, we extend (rather elegantly) any map \widehat{l}_n on $(sV)^{\wedge n}$ to a coderivation on the coalgebra $\Lambda(sV)$ by setting

$$\widehat{l}_n(sv_{[1 \text{ to } k]}) = \sum_{(n, k-n)} \widehat{\text{Kid}}(\rho) \widehat{l}_n(sv_{\rho[1 \text{ to } n]}) \wedge sv_{\rho[n+1 \text{ to } k]}. \quad (6)$$

The reader is left to verify that extending a map \widehat{l}_n as in equation (6) produces a well-defined map on $\Lambda(sV)$.

Although for any action ρ_n , extending an ρ_n -invariant map \widehat{l}_n as a coderivation on $T^c(sV)$ does not result in a ρ -invariant map on the tensor coalgebra, the restriction of the coderivation \widehat{l}_n on $T^c(sV)$ to the ρ -invariant subcoalgebra is equal to the coderivation one produces by extending \widehat{l}_n restricted to the ρ_n -invariant subspace of $(sV)^n$ as a coderivation on the ρ -invariant subcoalgebra of $T^c(sV)$. The set of all coderivations \widehat{l}_n generate the Lie algebra $\text{Coder}(\Lambda(sV))$. Next, we examine maps $\widehat{m}_n : (sV)^{n-1} \rightarrow W$ (the Lie action map \widehat{m}_2 is such a map) and $\widehat{f}_n : (sV)^n \rightarrow W$ (elements of $\text{Alt}_k(\mathfrak{g}, \mathbf{B})$ are of this form), where W is a graded commutative algebra. We will denote the degree of any element $w \in W$ by $|w|$. Elements of W have no external degree, at least not with respect to $T^c(sV)$ or its subcoalgebra $\Lambda(sV)$.

Subordinate coderivations and $\text{Coder}(\Lambda(sV) \rightarrow W)$

We begin with maps $\widehat{m}_n : (sV)^{n-1} \rightarrow W$. Let $\widehat{l}_n : (sV)^n \rightarrow sV$ be any map of degree $\text{id}(\widehat{l}_n)$ and let \widehat{l} be any coderivation on $T^c(sV)$ of degree $\text{id}(\widehat{l})$.

Definition 2.2. A map $\widehat{m}_n : (sV)^{n-1} \rightarrow W$ is an \widehat{l}_n -subordinate map if the degree of \widehat{m}_n and \widehat{l}_n agree. A map $\widehat{m} : T^c(sV) \rightarrow W$ is an \widehat{l} -subordinate coderivation if $\text{id}(\widehat{m}) = \text{id}(\widehat{l})$ and, for all p and q ,

$$(\Delta_{p,q} \rightarrow 1_W) \widehat{m} = (\widehat{l} \rightarrow 1)(\Delta_{p+n-1,q} \rightarrow 1_W) + (1 \rightarrow \widehat{m})(\Delta_{p,q+n-1} \rightarrow 1_W).$$

Just as the map \widehat{l}_n extends to a coderivation on $T^c(sV)$, so too, the map \widehat{m}_n may be extended as an \widehat{l}_n -subordinate coderivation. On $(sV)^k \rightarrow W$, we set $\widehat{m}_n = 0$ for $k < n-1$ and we set $\widehat{m}_n = \widehat{l}_n \rightarrow 1_W + 1 \rightarrow k-n+1 \rightarrow \widehat{m}_n$ for all $k > n-1$.

Definition 2.3. The map \widehat{m}_n is ρ_{n-1} -invariant if, for all $\sigma \in \Sigma_{n-1}$,

$$\widehat{m}_n(\sigma \cdot (sv_{[1 \text{ to } n-1]} \rightarrow w)) = \widehat{m}_n(sv_{[1 \text{ to } n-1]} \rightarrow w).$$

If both \widehat{l}_n and a subordinate coderivation \widehat{m}_n are invariant with respect to the same action ρ , then we may pass to the ρ -invariant subcoalgebra, where \widehat{m}_n remains subordinate to \widehat{l}_n . We extend the map \widehat{m}_n on $(sV)^{\wedge n-1} \rightarrow W$ as a coderivation on $\Lambda(sV) \rightarrow W$ by setting

$$\begin{aligned} \widehat{m}_n(sv_{[1 \text{ to } k]} \rightarrow w) = & \widehat{l}_n(sv_{\rho[1 \text{ to } k]} \rightarrow w) + \\ & \sum_{(k-n+1, n-1)} \widehat{\text{Kid}}(\sigma) \widehat{\text{Kid}}(\widehat{m}_n; sv_{\sigma[1 \text{ to } k-n+1]}) \\ & sv_{\sigma[1 \text{ to } k-n+1]} \widehat{m}_n(sv_{\sigma[k-n+2 \text{ to } k]} \rightarrow w). \end{aligned} \quad (7)$$

The set of all subordinate coderivations $\text{Coder}(\Lambda(sV) \rightarrow W)$ forms a graded Lie algebra under the graded commutator bracket. The bracket respects subordination, that is to say, if \widehat{m}_i and \widehat{m}_j are \widehat{l}_i and \widehat{l}_j -subordinate coderivations respectively, then $[\widehat{m}_i, \widehat{m}_j]$ is a coderivation subordinate to $[\widehat{l}_i, \widehat{l}_j]$. (Of course, the suspension map S respects subordinate maps and coderivations.)

The graded commutative algebra $\text{Hom}(\wedge(sV), \wedge(sV) \quad W)$

Suppose the map $\widehat{f}_n : (sV)^{\wedge n} \rightarrow W$ is suspended internal graded symmetric. We can extend \widehat{f}_n to an internal graded symmetric function from $\wedge(sV)$ into $\wedge(sV) \quad W$ by setting $\widehat{f}_n = 0$ for $k < n$ and, for $k \geq n$, setting $\widehat{f}_n(sv_{[1 \text{ to } k]}) =$

$$\sum_{\substack{\sigma \\ (k-n, n)}} \widehat{\text{K}}_{\text{id}}(\sigma) \widehat{\text{K}}_{\text{id}}(\widehat{f}_n; sv_{\sigma[1 \text{ to } k-n]}) sv_{\sigma[1 \text{ to } k-n]} \widehat{f}_n(sv_{\sigma[k-n+1 \text{ to } k]}). \quad (8)$$

The set of all such maps is $\text{Hom}(\wedge(sV), \wedge(sV) \quad W)$.

Suppose $\widehat{m}_p \in \text{Coder}(\wedge(sV) \quad W)$ is subordinate to $\widehat{l}_p \in \text{Coder}(\wedge(sV))$. The map

$$\langle \widehat{m}_p, \cdot \rangle : \text{Hom}(\wedge(sV), \wedge(sV) \quad W) \rightarrow \text{Hom}(\wedge(sV), \wedge(sV) \quad W),$$

given by setting

$$\langle \widehat{m}_p, \widehat{f}_n \rangle = \widehat{m}_p \widehat{f}_n - \widehat{\text{K}}_{\text{id}}(\widehat{m}_p; \widehat{f}_n) \widehat{f}_n \widehat{l}_p$$

on $(sV)^{\wedge n+p-1}$ and extending $\langle \widehat{m}_p, \widehat{f}_n \rangle$ to all of $\wedge(sV)$ as in equation (8) is well-defined. This map exposes an unusual type of graded Lie module structure on $\text{Hom}(\wedge(sV), \wedge(sV) \quad W)$. It is, in a sense, a Lie-module over *both* $\text{Coder}(\wedge(sV) \quad W)$ and $\text{Coder}(\wedge(sV))$ simultaneously in the sense that

$$\langle [\widehat{m}_i, \widehat{m}_j], \cdot \rangle = \langle \widehat{m}_i, \langle \widehat{m}_j, \cdot \rangle \rangle - \widehat{\text{K}}_{\text{id}}(\widehat{m}_i; \widehat{m}_j) \langle \widehat{m}_j, \langle \widehat{m}_i, \cdot \rangle \rangle.$$

We will call such a Lie module *shared*.

When (\mathbf{B}, sg) is a Lie-Rinehart pair, the Lie module map $\widehat{m}_2 : \mathfrak{g} \quad \mathbf{B} \rightarrow \mathbf{B}$ should be a map from sg into $\text{Der}(\mathbf{B})$. With this in mind, we introduce the following notion.

Definition 2.4. : An \widehat{l}_n -subordinate map \widehat{m}_n is a W -derivation source if for every $sv_{[1 \text{ to } n-1]}$ in $(sV)^{\wedge n-1}$, the map $\widehat{m}_n(sv_{[1 \text{ to } n-1]} \quad (\quad)) : W \rightarrow W$ is a graded derivation on W .

In other words, a W -derivation source \widehat{m}_n is a map from $(sV)^{\wedge n-1}$ into $\text{Der}(W)$. In practice, we will work with \widehat{m}_n as a coderivation on $\wedge(sV) \quad W$, where the characterization of \widehat{m}_n as a map into $\text{Der}(W)$ is somewhat obscured. The subset of all W -derivation sources in $\text{Coder}(\wedge(sV) \quad W)$ forms a graded Lie subalgebra denoted by $\text{Coder}^W(\wedge(sV) \quad W)$.

The subset of all coderivations $\widehat{l}_n \in \text{Coder}(\wedge(sV))$ which admit a subordinate W -derivation source \widehat{m}_n forms a graded Lie subalgebra which we will denote by $\text{Coder}^W(\wedge(sV))$.

The algebra structure on W extends to a graded commutative algebra structure on $\wedge(sV) \quad W$, which provides $\text{Hom}(\wedge(sV), \wedge(sV) \quad W)$ with a cup product. If \widehat{f}_n and \widehat{g}_s are maps in $\text{Hom}(\wedge(sV), \wedge(sV) \quad W)$. Then $\widehat{f}_n \smile \widehat{g}_s$ is given by setting $(\widehat{f}_n \smile \widehat{g}_s)(sv_{[1 \text{ to } n+s]}) =$

$$\sum_{\substack{\rho \\ (n, s)}} \widehat{\text{K}}_{\text{id}}(\rho) \widehat{\text{K}}_{\text{id}}(\widehat{g}_s; sv_{\rho[1 \text{ to } n]}) \widehat{f}_n(sv_{\rho[1 \text{ to } n]}) \cdot \widehat{g}_s(sv_{\rho[n+1 \text{ to } n+s]})$$

on $(sV)^{\wedge n+s}$, where \cdot represents the multiplication on W (henceforth, the \cdot will be suppressed). The map $\widehat{f}_n \smile \widehat{g}_s$ is then extended to all of $\wedge(sV)$ as in equation (8). The map $\langle \widehat{m}_p, \cdot \rangle$ acts as a derivation with respect to the cup product, so long as \widehat{m}_p is a W -derivation source.

W -linear maps and resting coderivations

For a Lie-Rinehart pair (\mathbf{B}, sg) , the Rinehart complex of \mathbf{B} -linear alternating maps is a subcomplex of the Chevalley-Eilenberg complex without modifying the differential. This is due to the \mathbf{B} -linearity of any \widehat{f}_n in $\text{Alt}_{\mathbf{B}}(\text{sg}, \mathbf{B})$, the Lie-Rinehart relations and the fact that the Lie module map \widehat{m}_2 is a \mathbf{B} -derivation source.

If the graded module V is a module over the graded commutative algebra W , then the tensor coalgebras $T^c V$ and $T^c(sV)$ are modules over TW . We will follow our general convention and denote $w_1 v_1 \cdots w_n v_n$ by $wv_{[1 \text{ to } n]}$ and, in the suspended setting, denote $w_1 s v_1 \cdots w_n s v_n$ by $wsv_{[1 \text{ to } n]}$. The *unraveling map*

$$\begin{aligned} \widehat{U} : T^c(sV) &\rightarrow W \otimes T^c(sV) \\ wsv_{[1 \text{ to } n]} &\mapsto \widehat{u}([1 \text{ to } n])w_{[1 \text{ to } n]} \otimes sv_{[1 \text{ to } n]}, \end{aligned}$$

where $\widehat{u}([1 \text{ to } n])$ is the sign produced when moving the w_i 's past the v_j 's. The map \widehat{U} respects the coalgebra structure of $T^c(sV)$, that is to say, for all p and q , the diagram

$$\begin{array}{ccc} (sV)^p & \xrightarrow{\Delta_{p,q}} & (sV)^q \\ \uparrow & & \uparrow \\ (sV)^{p+q} & \xrightarrow{\widehat{U}} & W \otimes (sV)^{p+q} \end{array} \quad \begin{array}{ccc} (sV)^p & \xrightarrow{\Delta_{p,q}} & (sV)^q \\ \uparrow & & \uparrow \\ (sV)^{p+q} & \xrightarrow{\widehat{U}} & W \otimes (sV)^{p+q} \end{array}$$

commutes.

Definition 2.5. : A suspended internal graded symmetric map $\widehat{f}_n : (sV)^{\wedge n} \rightarrow W$ is W -linear if

$$\widehat{f}_n(wsv_{[1 \text{ to } n]}) = \widehat{u}([1 \text{ to } n]) \widehat{\text{Kid}}(\widehat{f}_n; w_{[1 \text{ to } n]})w_{[1 \text{ to } n]}\widehat{f}_n(sv_{[1 \text{ to } n]}).$$

Likewise, a suspended internal graded symmetric map $\widehat{m}_n : (sV)^{\wedge n-1} \rightarrow W$ is W -linear if

$$\begin{aligned} \widehat{m}_n(wsv_{[1 \text{ to } n-1]} \otimes w) &= \\ \widehat{u}([1 \text{ to } n-1]) \widehat{\text{Kid}}(\widehat{m}_n; w_{[1 \text{ to } n-1]})w_{[1 \text{ to } n-1]}\widehat{m}_n(sv_{[1 \text{ to } n-1]} \otimes w). \end{aligned}$$

For a map $\widehat{m}_n : (sV)^{\wedge n-1} \rightarrow W$, we define the V -extension map $\widehat{m}_n^e : (sV)^{\wedge n} \rightarrow W$ by setting

$$\widehat{m}_n^e(wsv_{[1 \text{ to } n]}) = \sum_{\substack{\rho \\ (n-1,1)}} \widehat{\text{Kid}}(\rho)\widehat{m}_n(wsv_{\rho[1 \text{ to } n-1]} \otimes w_{\rho(n)})sv_{\rho(n)}.$$

Definition 2.6. : Given $\widehat{m}_n \in \text{Coder}^W(\wedge(sV) \rightarrow W)$, an W -linear, W -derivation source which is subordinate to $\widehat{l}_n \in \text{Coder}^W(\wedge(sV))$, the map \widehat{l}_n is said to rest on \widehat{m}_n if $\widehat{l}_n(wsv_{[1 \text{ to } n]}) =$

$$\widehat{u}([1 \text{ to } n]) \widehat{\text{Kid}}(\widehat{l}_n; w_{[1 \text{ to } n]})w_{[1 \text{ to } n]}\widehat{l}_n(sv_{[1 \text{ to } n]}) + \widehat{m}_n^e(wsv_{[1 \text{ to } n]}).$$

The subset of $\text{Coder}^W(\wedge(sV))$ of all coderivations \widehat{l}_p which rest on W -linear W -derivation sources \widehat{m}_p forms a graded Lie subalgebra. This subalgebra is denoted by $\text{Coder}_W^W(\wedge(sV))$. Similarly, the subset of $\text{Coder}^W(\wedge(sV) \rightarrow W)$ consisting of all W -linear W -derivation sources \widehat{m}_p subordinate to maps \widehat{l}_p , which in turn rest on \widehat{m}_p , forms a graded Lie subalgebra, denoted by $\text{Coder}_W^W(\wedge(sV) \rightarrow W)$. Proposition 2.7 is crucial and stated below; its proof is in the appendix.

PROPOSITION 2.7. : Let $\widehat{f}_n : (sV)^{\wedge n} \rightarrow W$ be a W -linear map and suppose the map $\widehat{m}_p : (sV)^{\wedge p-1} \rightarrow W$ is a W -linear, W -derivation source subordinate to the map $\widehat{l}_p : (sV)^{\wedge p} \rightarrow W$, which in turn rests on \widehat{m}_p . Then $\langle \widehat{m}_p, \widehat{f}_n \rangle = \widehat{m}_p \widehat{f}_n - \widehat{\text{Kid}}(\widehat{m}_p; \widehat{f}_n)\widehat{f}_n \widehat{l}_p$ is W -linear.

Proof of Proposition 2.7. : First we compute $\widehat{m}_p \widehat{f}_n(wsv_{[1 \text{ to } p+n-1]})$. We break up the result into two sums. The first one, $(\text{Lin } \widehat{m}_p \widehat{f}_n)$, consists of those terms in which all the w_i 's can and

have been factored out. The remaining terms form the second sum $(\text{Nonlin } \widehat{m}_p \widehat{f}_n)$. The first sum is over all $(p-1, n)$ -unshuffles σ :

$$(\text{Lin } \widehat{m}_p \widehat{f}_n) = \sum_{\sigma} \pm w_{[1 \text{ to } p+n-1]} \widehat{m}_p(sv_{\sigma[1 \text{ to } p-1]}^{\wedge}) \widehat{f}_n(sv_{\sigma[p \text{ to } n+p-1]}^{\wedge}).$$

The second sum is actually a double sum over all $(p-1, n)$ -unshuffles σ and all $(1, n-1)$ -unshuffles γ acting on the second σ -hand:

$$(\text{Nonlin } \widehat{m}_p \widehat{f}_n) = \sum_{\sigma} \sum_{\gamma} \pm \widehat{m}_p(sv_{\sigma[1 \text{ to } p-1]}^{\wedge}) w_{\sigma[1 \text{ to } p-1]} \cdot w_{\gamma\sigma[p \text{ to } p+n-1]} \cdot \widehat{f}_n(sv_{\gamma\sigma[p \text{ to } n+p-1]}^{\wedge}).$$

We compute $\widehat{f}_n \widehat{l}_p(wsv_{[1 \text{ to } p+n-1]}^{\wedge})$ by again breaking it up into two sums: $(\text{Lin } \widehat{f}_n \widehat{l}_p)$, is a sum over all $(p, n-1)$ -unshuffles ρ , i.e.,

$$(\text{Lin } \widehat{f}_n \widehat{l}_p) = \sum_{\rho} \pm w_{[1 \text{ to } p+n-1]} \widehat{f}_n(\widehat{l}_p(sv_{\sigma[1 \text{ to } p]}^{\wedge}) \wedge sv_{\sigma[p+1 \text{ to } n+p-1]}^{\wedge})$$

and $(\text{Nonlin } \widehat{f}_n \widehat{l}_p)$ is again a double sum over all $(p, n-1)$ -unshuffles ρ and all $(n-1, 1)$ -unshuffles τ acting on the first ρ -hand. In other words,

$$(\text{Nonlin } \widehat{f}_n \widehat{l}_p) = \sum_{\rho} \sum_{\tau} \pm \widehat{m}_p(sv_{\tau\rho[1 \text{ to } p-1]}^{\wedge}) w_{\tau\rho[1 \text{ to } p]} \cdot w_{\rho[p+1 \text{ to } p+n-1]} \widehat{f}_n(sv_{\tau\rho(p)} \wedge sv_{\rho[p+1 \text{ to } n+p-1]}^{\wedge}).$$

Recognizing that $\|\widehat{m}_p \widehat{f}_n\| = \|\widehat{f}_n \widehat{l}_p\|$, it follows that

$$\left[(\text{Lin } \widehat{m}_p \widehat{f}_n) - \widehat{\kappa}(\widehat{m}_p; \widehat{f}_n)(\text{Lin } \widehat{f}_n \widehat{l}_p) \right] = \pm w_{[1 \text{ to } p+n-1]} \left\langle \widehat{m}_p, \widehat{f}_n \right\rangle (sv_{[1 \text{ to } p+n-1]}^{\wedge}).$$

Therefore, once we show that $\left[(\text{Nonlin } \widehat{m}_p \widehat{f}_n) - \widehat{\kappa}(\widehat{m}_p; \widehat{f}_n)(\text{Nonlin } \widehat{f}_n \widehat{l}_p) \right] = 0$, the proof will be complete.

For every pair consisting of a $(p-1, n)$ -unshuffle σ and a $(1, n-1)$ -unshuffle γ acting on the second σ hand, there is a unique pair consisting of a $(p, n-1)$ -unshuffle ρ and a $(p-1, 1)$ -unshuffle τ acting on the first ρ hand which produces the exact same sequence of components, that is to say,

$$\begin{aligned} \sigma[1 \text{ to } p-1] &= \tau\rho[1 \text{ to } p-1], \\ \gamma\sigma(p) &= \tau\rho(p) \text{ and} \\ \gamma\sigma[p+1 \text{ to } p+n-1] &= \rho[p+1 \text{ to } p+n-1]. \end{aligned}$$

Why should this be so? A simple counting argument provides the answer. Either combination produces an $(p-1, 1, n-1)$ -unshuffle. There are $\binom{p+n-2}{p-1}$ distinct $(p-1, 1, n-1)$ -unshuffles which send $j \in \{1, \dots, p+n-1\}$ to the p^{th} position because once the p^{th} position is determined, the remaining $p+n-2$ terms must be unshuffled into a hand of length $p-1$ and a hand of length $n-1$. How many distinct pairs (σ, γ) send j to the p^{th} position? This question is equivalent to asking how many distinct $(p-1, n)$ -unshuffles σ send j to the second hand since, once j is in the second hand, there is a unique $(1, n-1)$ unshuffle γ which will send j to the p^{th} position. The number of $(p-1, n)$ -unshuffles is $\binom{p+n-1}{n}$ and $\binom{p+n-2}{n-1}$ of them have j in the second hand. Since $\binom{p+n-2}{n-1} = \binom{p+n-2}{p-1}$, we know there is a one-to-one correspondence between $(p-1, 1, n-1)$ -unshuffles and pair (σ, γ) which send j to the p^{th} position. A similar argument shows that the number of $(p-1, 1, n-1)$ -unshuffles which send j to the p^{th} position equals the number of (ρ, τ) combinations which do the same. We can show that the (σ, γ) term of $(\text{Nonlin } \widehat{m}_p \widehat{f}_n)$ and the corresponding (ρ, τ) term of $(\text{Nonlin } \widehat{f}_n \widehat{l}_p)$ have opposite signs, so they cancel. \square

Naturally, the graded commutative algebra $\text{Hom}_W(\Lambda(sV), \Lambda(sV) \otimes W)$ is a shared Lie module over the Lie algebras $\text{Coder}_W^W(\Lambda(sV), \Lambda(sV) \otimes W)$ and $\text{Coder}_W^W(\Lambda(sV))$.

3. Lie algebras, their modules and cohomology in the coalgebra setting

Lie algebras and their modules

The most familiar definition of a Lie algebra \mathfrak{g} is as an ungraded k -vector space (or, more generally, a module over a k -algebra A) equipped with a skew-symmetric linear map $l_2 = [\ , \] : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ which satisfies the Jacobi identity, that is to say the equality

$$l_2(x, l_2(y, z)) = l_2(l_2(x, y), z) + l_2(y, l_2(x, z))$$

holds for all $x, y, z \in \mathfrak{g}$. In the context of graded coassociative coalgebras, we can consider \mathfrak{g} a graded vector space where all elements of \mathfrak{g} have combined degree $\langle 1, 0 \rangle$. The bracket l_2 becomes a degree $\langle -1, 0 \rangle$ coderivation on $\bigwedge \mathfrak{g}$. No information is lost in the process because

$$\begin{aligned} l_2^{\otimes} (x \otimes y) &= l_2 \left(x \otimes y + (-1)^{\langle 0,1 \rangle \cdot \langle 0,1 \rangle} y \otimes x \right) \\ &= l_2(x \otimes y) - l_2(y \otimes x) \\ &= l_2(x \otimes y) + l_2(x \otimes y) \\ &= 2l_2(x \otimes y). \end{aligned}$$

But we can also define the Lie algebra on $\bigwedge(\mathfrak{sg})$, where the bracket \widehat{l}_2 becomes an internal graded symmetric coderivation of degree -1 on $\bigwedge(\mathfrak{sg})$ with respect to the internal grading. Again, no essential information is lost. We can therefore realize a Lie algebra in three equivalent ways:

1. A Lie algebra is a vector space \mathfrak{g} over k with skew symmetric bracket l_2 which satisfies the Jacobi identity (the original definition).
2. A Lie algebra is the subcoalgebra $\bigwedge \mathfrak{g}$, together with a combined degree $\langle -1, 0 \rangle$ coderivation l_2 satisfying $l_2 \circ l_2 = \frac{1}{2}[l_2, l_2] = 0$, which is equivalent to the Jacobi identity. Since the internal degree is zero, it can be ignored, so l_2 can be considered a degree -1 coderivation, i.e., a differential on $\bigwedge \mathfrak{g}$.
3. A Lie algebra is a subcoalgebra $\bigwedge(\mathfrak{sg})$ together with a suspended internal degree -1 differential \widehat{l}_2 . This is our preferred setting.

The Jacobi identity is a result of the more general fact, true for all coderivations \widehat{l} with odd degree, that $[\widehat{l}, \widehat{l}] = 2\widehat{l} \circ \widehat{l}$ and $[\widehat{l}, \widehat{l}] = -[\widehat{l}, \widehat{l}] = 0$. In other words, any degree -1 coderivation \widehat{l} is a differential on the coalgebra $\bigwedge(\mathfrak{sg})$ and if the differential \widehat{l} has external degree -1 , then it is the extension of a map $\widehat{l} : \mathfrak{sg} \wedge \mathfrak{sg} \rightarrow \mathfrak{sg}$ as a coderivation, i.e., \widehat{l} is a Lie bracket.

Suppose \mathbf{B} is an ungraded Lie module over \mathfrak{g} , i.e., there is a map $\widehat{m}_2 : \mathfrak{sg} \otimes \mathbf{B} \rightarrow \mathbf{B}$ such that $\widehat{m}_2(\widehat{l}_2(sx \wedge sy) \otimes b) - \widehat{m}_2(sx \otimes \widehat{m}_2(sy \otimes b)) + \widehat{m}_2(sy \otimes \widehat{m}_2(sx \otimes b)) = 0$, for all $x, y \in \mathfrak{g}$ and $b \in \mathbf{B}$. The map \widehat{m}_2 is \widehat{l}_2 -subordinate, and we can extend \widehat{m}_2 to an \widehat{l}_2 -subordinate coderivation on $\bigwedge(\mathfrak{sg}) \otimes \mathbf{B}$. The equation above can be rewritten as $\widehat{m}_2 \circ \widehat{m}_2 = \frac{1}{2}[\widehat{m}_2, \widehat{m}_2] = 0$ and since the degree of \widehat{m}_2 is -1, it is a differential on $\bigwedge(\mathfrak{sg}) \otimes \mathbf{B}$. In the coalgebra setting, then, an \mathfrak{g} -module structure is any coderivation \widehat{m}_2 on $\bigwedge(\mathfrak{sg}) \otimes \mathbf{B}$ which is a differential and which is subordinate to the bracket \widehat{l}_2 .

(Cartan-)Chevalley-Eilenberg Cohomology

Definition 3.1. : A Chevalley-Eilenberg pair $(\mathbf{B}, \mathfrak{g})$ is a Lie algebra \mathfrak{g} and Lie module \mathbf{B} over \mathfrak{g} .

For any Chevalley-Eilenberg pair $(\mathbf{B}, \mathfrak{g})$, an n -multilinear function $f_n : \mathfrak{g}^{\times n} \rightarrow \mathbf{B}$ is *alternating* if

$$f_n(x_{\sigma(1)}, \dots, x_{\sigma(n)}) = (-1)^\sigma f_n(x_1, \dots, x_n)$$

for all $\sigma \in \Sigma_n$, where $(-1)^\sigma$ is the sign of the permutation. The algebra of all alternating multilinear functions $Alt_k(\mathfrak{g}, \mathbf{B})$ is graded by n and admits a degree +1 differential $\delta_{CE} : Alt_k^n(\mathfrak{g}, \mathbf{B}) \rightarrow Alt_k^{n+1}(\mathfrak{g}, \mathbf{B})$ given by equation (2). The cohomology of this complex with respect to δ_{CE} is the *Chevalley-Eilenberg cohomology of \mathfrak{g} with coefficients in \mathbf{B}* .

In the coalgebra setting, any alternating n -multilinear function f_n can be realized uniquely as a linear function $\widehat{f}_n : (\mathfrak{sg})^{\wedge n} \rightarrow \mathbf{B}$, which then can be extended as a coderivation $\widehat{f}_n : \wedge(\mathfrak{sg}) \rightarrow \wedge(\mathfrak{sg}) \quad \mathbf{B}$. We extend $b \in \mathbf{B}$ as a coderivation $b : \wedge(\mathfrak{sg}) \rightarrow \wedge(\mathfrak{sg}) \quad \mathbf{B}$ by setting $b(sx_{[1 \text{ to } k]}) = sx_{[1 \text{ to } k]} \quad b$. Any function $\widehat{F} \in \text{Hom}(\wedge(\mathfrak{sg}), \wedge(\mathfrak{sg}) \quad \mathbf{B})$ is the sum $\sum_{n=0}^\infty \widehat{f}_n$, where $\widehat{f}_n : (\mathfrak{sg})^{\wedge n} \rightarrow \mathbf{B}$ has degree n . Hence, $\text{Hom}(\wedge(\mathfrak{sg}), \wedge(\mathfrak{sg}) \quad \mathbf{B})$ is graded by \mathbb{N} and is isomorphic to $Alt_k(\mathfrak{g}, \mathbf{B})$. The differential δ_{CE} on $\text{Hom}(\wedge(\mathfrak{sg}), \wedge(\mathfrak{sg}) \quad \mathbf{B})$ has the elegant form $\langle \widehat{m}_2, \cdot \rangle$. This formula works even for $b \in \mathbf{B}$, since $\delta_{CE}b = \langle \widehat{m}_2, b \rangle = \widehat{m}_2b + b\widehat{l}_2$ and $\widehat{l}_2 = 0$ on \mathfrak{sg} . Since $[\widehat{m}_2, \widehat{m}_2] = -[\widehat{m}_2, \widehat{m}_2] = 0$, it is now simple to show that

$$\delta_{CE} \circ \delta_{CE} = \langle \widehat{m}_2 \langle \widehat{m}_2, \cdot \rangle \rangle = \left\langle \frac{1}{2} [\widehat{m}_2, \widehat{m}_2], \cdot \right\rangle = 0.$$

If \mathbf{B} is an algebra, then $\text{Hom}(\wedge(\mathfrak{sg}), \wedge(\mathfrak{sg}) \quad \mathbf{B})$ is also an algebra. The differential δ_{CE} acts as a derivation with respect to the multiplication.

Lie-Rinehart pairs and Rinehart cohomology

Suppose \mathbf{B} is an algebra over k which is also a \mathfrak{g} -module. The definition of a Lie-Rinehart pair below has been modified from definition 1.1 to reflect the coalgebra setting.

Definition 3.2. : [Rin63] Let $\mu : \mathbf{B} \quad (\mathfrak{sg}) \rightarrow (\mathfrak{sg})$ be a left \mathbf{B} -module action on \mathfrak{sg} denoted by $\mu(a \quad \alpha) := a\alpha$. Let $\widehat{m}_2 : (\mathfrak{sg}) \quad \mathbf{B} \rightarrow \mathbf{B}$ be an \widehat{l}_2 -subordinate \mathbf{B} -derivation source, i.e., an \mathfrak{sg} -module action on \mathbf{B} . The pair $(\mathbf{B}, \mathfrak{sg})$ is a Lie-Rinehart pair, provided the Lie-Rinehart relations (LRa) and (LRb) are satisfied.

LRa: \widehat{m}_2 is \mathbf{B} -linear.

LRb: \widehat{l}_2 rests on \widehat{m}_2 (definition 2.6).

Suppose \widehat{f}_n is in $\text{Hom}_{\mathbf{B}}(\wedge(\mathfrak{sg}), \wedge(\mathfrak{sg}) \quad \mathbf{B})$. From proposition 2.7, we know the image of \widehat{f}_n under the Chevalley-Eilenberg differential $\delta_{CE} = \langle \widehat{m}_2, \cdot \rangle$ is again \mathbf{B} -linear. The Rinehart algebra $R = \text{Hom}_{\mathbf{B}}(\wedge(\mathfrak{sg}), \wedge(\mathfrak{sg}) \quad \mathbf{B})$ with differential $\delta_R = \langle \widehat{m}_2, \cdot \rangle$ is a subcomplex of the Chevalley-Eilenberg complex and the cohomology of R with respect to δ_R is the *Rinehart cohomology of \mathfrak{g} with coefficients in \mathbf{B}* .

4. Strongly homotopy Lie algebras, their modules and cohomology

Strongly homotopy Lie algebras (shLie algebras) first appeared implicitly in [Sul77] and explicitly in [SS] in the context of deformation theory. A concise introduction to shLie algebras is found in [LM95]. The initial definitions below are lifted directly from [LM95] and then modified to fit the language introduced in §2.

Definition 4.1. : [LM95] and [LS] An $\mathcal{L}(m)$ -structure on a graded module L is a system of linear maps $\{l_k : 1 \leq k \leq m \leq \infty, k \neq \infty\}$, where $l_k : \otimes^k L \rightarrow L$ has combined degree $\langle 1 - k, k - 2 \rangle$ and each map is combined graded symmetric in the sense that

$$l_k(v_{\sigma[1 \text{ to } k]}) = \mathbf{K}_{||} ||(\sigma)l_k(v_{[1 \text{ to } k]})$$

for all $\sigma \in \Sigma_k$. Moreover, the following generalized form of the Jacobi identity, the n^{th} Jacobi identity map, is satisfied for $n \leq m$:

$$\mathcal{J}_{\text{ID}_n}(v_{[1 \text{ to } n]}) = \sum_{\substack{i+j=n+1 \\ i,j \geq 1}} \sum_{\sigma} \mathbb{K}_{||}(\sigma)(-1)^{i(j-1)} l_j(l_i(v_{\sigma(1 \text{ to } i)})) \quad v_{\sigma(i+1 \text{ to } n)} = 0,$$

where the second summation runs over all $(i, j - 1)$ -unshuffles. The graded vector space L is a strongly homotopy Lie algebra (or shLie algebra) if L admits an $\mathcal{L}(\infty)$ -structure.

We can rewrite this definition in terms of maps on $\bigwedge L$ because the maps l_k are combined graded symmetric. The Jacobi identity maps $\mathcal{J}_{\text{ID}_n}$ should equal 0 on $\bigwedge L$, but it is not immediately evident that the maps $\mathcal{J}_{\text{ID}_n}$ can be extended as coderivations on $\bigwedge L$. Fortunately, we can rewrite the Jacobi identity maps using the bracket on $\text{Coder}(\bigwedge L)$. Consider the bracket of l_j with l_i with respect to the combined grading on $\bigwedge L$:

$$[l_j, l_i] = l_j l_i - (-1)^{(1-j, j-2) \cdot (1-i, i-2)} l_i l_j = l_j l_i + (-1)^{j+i} l_i l_j.$$

Multiplying both sides by the desired sign for $l_j l_i$ in the Jacobi identity map $\mathcal{J}_{\text{ID}_n}$, we find that

$$\begin{aligned} (-1)^{i(j-1)} [l_j, l_i] &= (-1)^{i(j-1)} l_j l_i + (-1)^{i(j-1)+j+i} l_i l_j \\ &= (-1)^{i(j-1)} l_j l_i + (-1)^{j(i-1)} l_i l_j. \end{aligned}$$

We therefore reformulate the definition of an $\mathcal{L}(m)$ -algebra and an shLie algebra:

Definition 4.2. : An $\mathcal{L}(m)$ -structure on a graded module L consists of a system of coderivations $\{l_k : \bigwedge L \rightarrow \bigwedge L : 1 \leq k \leq m \leq \infty, k \neq \infty\}$, where each l_k has combined degree $\langle 1 - k, k - 2 \rangle$. Moreover, the following generalized form of the Jacobi identity is satisfied for $n \leq m$:

$$\mathcal{J}_{\text{ID}_n} = \frac{1}{2} \sum_{\substack{i+j=n+1 \\ i,j \geq 1}} (-1)^{i(j-1)} [l_j, l_i] = 0 \tag{9}$$

on $L^{\otimes n}$. Again, if L admits an $\mathcal{L}(\infty)$ -structure, then L is a strongly homotopy Lie algebra (an shLie algebra).

Since the Jacobi identity maps are sums of coderivations, it follows that $\mathcal{J}_{\text{ID}_n}$ can be extended to a coderivation on $\bigwedge L$. Furthermore, since $\mathcal{J}_{\text{ID}_n} = 0$ on $L^{\otimes n}$, we have proven the following proposition.

PROPOSITION 4.3. : $\mathcal{J}_{\text{ID}_n} = 0$ on $\bigwedge L$.

In both [LS] and [LM95], this result was not achieved without first passing to the suspended internal graded symmetric setting. Lada, Markl and Stasheff defined strongly homotopy Lie algebras on the tensor coalgebra $T^c L$ rather than on $\bigwedge L$, whose existence had not been recognized. The maps $\mathcal{J}_{\text{ID}_n}$ cannot be extended as coderivations on the tensor coalgebra while preserving the desired symmetry. So instead, they suspended the maps l_k , changing the symmetry of the maps from combined graded symmetric to suspended internal graded symmetric. They were then in familiar territory, where the maps $\widehat{\mathcal{J}}_{\text{ID}_n}$ could be extended as coderivations on $\widehat{\bigwedge}(sL)$, even though they did not make explicit use of the bracket on $\text{Coder}(\widehat{\bigwedge}(sL))$.

Modules over a strongly homotopy Lie algebra

Definition 4.4. : [LM95] Let $L = (L, l_i)$ be an $\mathcal{L}(p)$ -algebra ($0 < p < \infty$) and let M be a differential graded module with differential m_1 . Then a left $\mathcal{L}(k)$ -module structure over L on M (for $k \leq p$) is a collection $\{m_n : 1 \leq n \leq k, n \neq \infty\}$ of l_n -subordinate coderivations m_n on $\bigotimes L \rightarrow M$ such that the n^{th} action identity map

$$\begin{aligned} ACT_{\text{ID}_n} &= \sum_{i+j=n+1} (-1)^{i(j-1)} m_j m_i \\ &= \frac{1}{2} \sum_{\substack{i+j=n+1 \\ i,j \geq 1}} (-1)^{i(j-1)} [m_j, m_i] \\ &= 0 \end{aligned}$$

on $\bigotimes L \rightarrow M$. The differential graded module M is a strongly homotopy Lie module over L (or an L -shLie module) if L admits an $\mathcal{L}(\infty)$ -structure and M is a module with respect to that $\mathcal{L}(\infty)$ -structure.

Definition 4.4 implies that the differential m_1 on M must be l_1 -subordinate. It is simple to verify that m_1 is a differential on $\bigotimes L \rightarrow M$.

When M is a differential graded commutative algebra, we require that the maps m_i be M -derivation sources, that is to say, for every $v_{[1 \text{ to } i-1]} \in L^{\otimes i-1}$, the map $m_i(v_{[1 \text{ to } i-1]} \quad (\quad)) : M \rightarrow M$ is in $\text{Der}(M)$.

Since $\text{Coder}(\bigotimes L) \approx \text{Coder}(\wedge(sL))$, it follows that if (L, l_i) is a strongly homotopy Lie algebra, then (sL, \widehat{l}_i) , where \widehat{l}_i is the coderivation on $\wedge(sL)$ induced by the isomorphism $S : \bigotimes L \rightarrow \wedge(sL)$, is also an shLie algebra. The suspended internal degree of each map \widehat{l}_i is -1 , so the sum of the $\mathcal{L}(\infty)$ -structure maps \widehat{l}_i is a differential on $\wedge(sL)$, first shown in [LS]. This is not the case for (L, l_i) because each l_i is bigraded. In order for the sum of the l_i 's to be a differential, one would have to make sense of l_i as a degree -1 map. Passing to the suspended internal graded symmetric setting is a convenient way to accomplish this because degree of \widehat{l}_i is $\text{ed}(l_i) + \text{id}(l_i) = -1$, which is the total degree of l_i . For (sL, \widehat{l}_i) , the n^{th} Jacobi identity map is

$$\widehat{\mathcal{J}}_{\text{ID}_n} = \frac{1}{2} \sum_{\substack{i+j=n+1 \\ i,j \geq 1}} [\widehat{l}_j, \widehat{l}_i], \tag{10}$$

which is 0 on $\wedge(sL)$.

PROPOSITION 4.5. : [LS] If (sL, \widehat{l}_i) is an shLie algebra, the map $D_{sL} = \sum_{i=1}^{\infty} \widehat{l}_i$ is a differential on $\wedge(sL)$, i.e. it is a map of degree -1 such that $D_{sL} \circ D_{sL} = 0$.

Proof of Proposition 4.13. : A proof can be found in [LS]. Here the proof is essentially the same but takes advantage of the bracket of coderivations:

$$D_{sL} \circ D_{sL} = \left(\sum_{j=1}^{\infty} \widehat{l}_j \right) \circ \left(\sum_{i=1}^{\infty} \widehat{l}_i \right) = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} \widehat{l}_j \widehat{l}_i = \sum_{n=1}^{\infty} \sum_{\substack{i+j=n+1 \\ i,j \geq 1}} \widehat{l}_j \widehat{l}_i.$$

Using equation (10) in definition 4.2, we see that

$$\sum_{n=1}^{\infty} \sum_{\substack{i+j=n+1 \\ i,j \geq 1}} \widehat{l}_j \widehat{l}_i = \frac{1}{2} \sum_{n=1}^{\infty} \sum_{\substack{i+j=n+1 \\ i,j \geq 1}} [\widehat{l}_j, \widehat{l}_i] = \frac{1}{2} \sum_{n=1}^{\infty} \widehat{\mathcal{J}}_{\text{ID}_n} = 0.$$

Note also that D_{sL} is a degree -1 coderivation, so $D_{sL} \circ D_{sL} = \frac{1}{2} [D_{sL}, D_{sL}]$ and the graded skew-commutativity of the bracket ensures that $[D_{sL}, D_{sL}] = 0$. \square

Suppose (M, m_i) is an shLie module over (L, l_i) . In the suspended internal graded symmetric setting, the maps $\widehat{m}_i : \wedge(sL) \rightarrow M$ form an shLie module structure on M over (sL, \widehat{l}_i) . Again, the degree of \widehat{m}_i is -1 , the total degree of m_i , and the action identity map has the form

$$\widehat{ACT}_{\text{ID}_n} = \frac{1}{2} \sum_{\substack{i+j=n+1 \\ i, j \geq 1}} [\widehat{m}_j, \widehat{m}_i], \quad (11)$$

which is zero on $\wedge(sL) \rightarrow M$.

PROPOSITION 4.6. : *The map $D_M = \sum_{i=1}^{\infty} \widehat{m}_i$ is a differential on $\wedge(sL) \rightarrow M$, i.e., $D_M \circ D_M = 0$.*

Proof of Proposition 4.14. : Isomorphic to the proof of proposition 4.5. Notice that D_M is D_{sL} -subordinate and that $D_M \circ D_M = \frac{1}{2}[D_M, D_M] = 0$. \square

It is only for historic reasons that we present shLie structures and shLie module structures first in the combined graded symmetric setting first and then in the suspended internal graded symmetric one. We will now use the suspended internal graded symmetric setting exclusively.

Homotopy Chevalley-Eilenberg cohomology

Definition 4.7. : *A homotopy Chevalley-Eilenberg pair (M, L) consists of an shLie algebra L and an L -shLie module M .*

A multi-linear alternating function $F_n : L^{\times n} \rightarrow M$ can be seen as a linear function $\widehat{F}_n : (sL)^{\wedge n} \rightarrow M$. So the homotopy Chevalley-Eilenberg algebra $\text{Alt}_k(L, M)$ is isomorphic to

$$\text{Hom}_k(\wedge(sL), \wedge(sL) \rightarrow M).$$

Any homogeneous map \widehat{F} splits into a sum $\sum_{n=0}^{\infty} \widehat{F}_n$, where $\widehat{F}_n : (sL)^{\wedge n} \rightarrow M$. The map \widehat{F}_n is extended as a coderivation on $\wedge(sL) \rightarrow M$. Any function \widehat{F}_0 is an element $a \in M$. The degree of $a : (sL)^{\wedge 0} \rightarrow M$ as a map is the negative of its degree as an element of M . The map a is extended as a coderivation on $\wedge(sL) \rightarrow M$ by setting

$$a(sv_{[1 \text{ to } k]}^{\wedge}) = \widehat{\mathbf{K}}_{\text{id}}(a; sv_{[1 \text{ to } k]}^{\wedge})sv_{[1 \text{ to } k]}^{\wedge} \quad a.$$

Instead of \mathbb{N} -graded, as was the case in the ungraded setting, the *Chevalley-Eilenberg complex* $\text{Hom}_k(\wedge(sL), \wedge(sL) \rightarrow M)$ is \mathbb{Z} -graded by suspended internal degree. For any $\widehat{F} \in \text{Hom}_k(\wedge(sL), \wedge(sL) \rightarrow M)$ of degree $\widehat{\text{id}}(\widehat{F})$, we define

$$\delta_{hCE}\widehat{F} = D_M\widehat{F} - \widehat{\mathbf{K}}_{\text{id}}(\widehat{F})\widehat{F}D_{sL} = \langle D_M, \widehat{F} \rangle,$$

which has degree $\widehat{\text{id}}(\widehat{F}) + 1$.

PROPOSITION 4.8. : $\delta_{hCE} \circ \delta_{hCE} = 0$.

Proof of Proposition 4.17. :

$$\begin{aligned} \delta_{hCE} \circ \delta_{hCE} &= \langle D_M, \langle D_M, \widehat{F} \rangle \rangle \\ &= \left\langle \frac{1}{2}[D_M, D_M], \widehat{F} \right\rangle \\ &= 0. \quad \square \end{aligned}$$

The cohomology with respect to δ_{hCE} is the *homotopy Chevalley-Eilenberg cohomology* of sL with coefficients in M .

When M is a differential graded commutative algebra, the homotopy Chevalley-Eilenberg complex $\text{Hom}_k(\wedge(sL), \wedge(sL) \otimes M)$ is a differential graded commutative algebra. Suppose $\widehat{F} = \sum_{i=0}^{\infty} \widehat{F}_i$ and $\widehat{G} = \sum_{j=0}^{\infty} \widehat{G}_j$ are both elements of the complex, where $\widehat{F}_i : \wedge^i(sL) \rightarrow M$ and $\widehat{G}_j : \wedge^j(sL) \rightarrow M$. The product $\widehat{F} \smile \widehat{G} = \sum_{n=0}^{\infty} (\widehat{F} \smile \widehat{G})_n$ is given by $(\widehat{F} \smile \widehat{G})_n = \sum_{i+j=n} \widehat{F}_i \smile \widehat{G}_j$. The differential δ_{hCE} acts as a derivation with respect to this multiplication.

Homotopy Lie-Rinehart pairs and homotopy Rinehart cohomology

The *homotopy Rinehart complex* is a straightforward generalization of the Rinehart complex in the ungraded setting. Since the Rinehart complex is defined only for Lie-Rinehart pairs $(\mathbf{B}, \mathfrak{g})$, we must define what constitutes a homotopy Lie-Rinehart pair (M, sL) . The subset $\text{Alt}_M(sL, M)$ of $\text{Alt}_k(sL, M)$ consisting of all M -linear alternating functions is isomorphic to $\text{Hom}_M(\wedge(sL), \wedge(sL) \otimes M)$.

Definition 4.9. : A homotopy Lie-Rinehart pair (M, sL) consists of a differential graded commutative algebra (M, \widehat{m}_i) which is an shLie module over the shLie algebra (sL, \widehat{l}_i) , which in turn is an M -module. Moreover, the following two homotopy Lie-Rinehart relations must be satisfied for all $i \geq 1$:

- (hLRa_i): The shLie module structure map \widehat{m}_i is M -linear.
- (hLRb_i): The shLie structure map \widehat{l}_i rests on \widehat{m}_i (definition 2.6).

An $\mathcal{L}(p)$ -Lie-Rinehart pair $((M, \widehat{m}_i), (sL, \widehat{l}_i))$ has maps \widehat{m}_i and \widehat{l}_i which satisfy (hLRa_i) and (hLRb_i) for $1 \leq i \leq p$.

PROPOSITION 4.10. : If \widehat{F} is M -linear, then so is $\delta_{hCE}\widehat{F}$.

Proof of Proposition 4.18. : The image of \widehat{F} under δ_{hCE} is

$$\langle D_M, \widehat{F} \rangle = \sum_{i=1}^{\infty} \langle \widehat{m}_i, \widehat{F} \rangle.$$

The proof follows as a consequence of proposition 2.7. □

We conclude that the subset of all M -linear functions in $\text{Hom}_k(\wedge(sL), \wedge(sL) \otimes M)$ forms a subcomplex $\mathcal{R} = \text{Hom}_M(\wedge(sL), \wedge(sL) \otimes M)$ with differential $\delta_{\mathcal{R}} = \langle D_M, \cdot \rangle$. The differential, together with the cup product, provides the homotopy Rinehart complex \mathcal{R} with the structure of a differential graded commutative algebra. The cohomology of \mathcal{R} with respect to $\delta_{\mathcal{R}}$ is the *homotopy Rinehart cohomology* of sL with coefficients in M .

5. Homotopy Lie-Rinehart resolutions of Lie-Rinehart pairs

It is often necessary to replace the components of a complex with resolutions of those component in order to construct a *model* for a complex with the same basic algebraic structure and precisely the same (co)homology as the original. (It is not always necessary to replace all of the components, see [Sta92], for example.) When building a model for the Rinehart cohomology for a Lie-Rinehart pair $(\mathbf{B}, \mathfrak{g})$ over a k -algebra A , we replace the pair with a pair of resolutions (M, sL) , which retain as much of the algebraic structure of $(\mathbf{B}, \mathfrak{g})$ as possible. It makes sense, then, that (M, L) should form a homotopy Lie-Rinehart pair.

Piecing together Lie-Rinehart resolutions

Let $(\mathbf{B}, \mathfrak{g})$ be a Lie-Rinehart pair over a k -algebra A with module structure maps $\widehat{\mu} : \mathbf{B} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ and $\widehat{m}_2 : \mathfrak{g} \otimes \mathbf{B} \rightarrow \mathbf{B}$. Let \widehat{l}_2 denote the bracket on \mathfrak{g} and $\pi_{\mathbf{B}}$ denote the multiplication on \mathbf{B} . (Henceforth, only maps associated with the Lie-Rinehart pair will have hats.) The basic ingredients of a homotopy Lie-Rinehart resolution for a Lie-Rinehart pair $(\mathbf{B}, \mathfrak{g})$ are (sL, l_i) and (M, m_i, π_M) where $H_{l_1}(sL) = \mathfrak{g}$ and $H_{m_1}(M) = \mathbf{B}$.

Definition 5.1. : A homotopy Lie-Rinehart resolution of a Lie-Rinehart pair $(\mathbf{B}, \mathbf{sg})$ over an algebra A is a homotopy Lie-Rinehart pair (M, sL) over A , such that

(a) the shLie algebra (sL, l_i) , seen as the coalgebra $\bigwedge(sL)$, resolves $\bigwedge(\mathbf{sg})$, i.e.,

$$H_{l_1}(\bigwedge(sL)) = \bigwedge(\mathbf{sg}), \tag{12}$$

where (following the physicists' notation) H_{l_1} denotes the homology with respect to the differential l_1 ,

(b) the differential graded commutative algebra (M, m_i, π_M) satisfies

$$H_{m_1}(\bigwedge(sL) \quad M) = \bigwedge(\mathbf{sg}) \quad \mathbf{B}, \tag{13}$$

(c) the dgca (M, m_i, π_M) also satisfies

$$H_{m_1}(M \quad M) = \mathbf{B} \quad \mathbf{B}. \tag{14}$$

Furthermore, the following conditions hold:

- i. $H_{l_1}(l_2) = \widehat{l_2}$ on $\bigwedge(\mathbf{sg})$.
- ii. $H_{m_1}(m_2) = \widehat{m_2}$ on $\bigwedge(\mathbf{sg}) \quad \mathbf{B}$.
- iii. $H_{m_1}(\pi_M) = \pi_{\mathbf{B}}$ on $\mathbf{B} \quad \mathbf{B}$.

The conditions above are quite reasonable, but satisfying them is far from automatic. Consider any homotopy Lie-Rinehart pair (M, sL) . Since m_1 is a derivation on M , the multiplication on M is a chain map which induces a well-defined map $H_{m_1}(\pi_M)$ from $H_{m_1}(M \quad M)$ into $H_{m_1}(M)$. Both $\bigwedge(sL)$ and $\bigwedge(sL) \quad M$ are complexes with differentials l_1 and m_1 , respectively. Since the maps $\mathcal{J}_{\text{ID}_2}$ and $\mathcal{ACT}_{\text{ID}_2}$ are both zero, it follows that both l_2 and m_2 are graded morphisms of complexes and as such, induce well-defined maps $H_{l_1}(l_2)$ and $H_{m_1}(m_2)$ on their respective homologies.

In the proposition below, we state conditions on the homotopy Lie-Rinehart pair which guarantees it satisfies the conditions in definition 5.1.

PROPOSITION 5.2. : Let (M, sL) be a homotopy Lie-Rinehart pair. Suppose the differential graded algebra (M, m_i, π_M) is a projective resolution of \mathbf{B} over A which respects the algebra structure on \mathbf{B} , i.e., condition iii of definition 5.1 is satisfied. Suppose (sL, l_i) is a projective resolution \mathbf{sg} . Furthermore,

- (a) $H_{l_1}(l_2) = \widehat{l_2}$ on $\mathbf{sg} \wedge \mathbf{sg}$ and
- (b) $H_{m_1}(m_2) = \widehat{m_2}$ on $\mathbf{sg} \quad \mathbf{B}$.

Then (M, sL) is a homotopy Lie-Rinehart resolution of $(\mathbf{B}, \mathbf{sg})$.

SDR-Data

For the proof of proposition 5.2, we will use strong deformation retract data (SDR-data). In general, SDR-data for two differential graded modules (M, d_M) and (N, d_N) over a commutative ring R with unity consist of two degree 0 chain maps $\lambda : M \rightarrow N$ and $p : N \rightarrow M$ and a homotopy $h : N \rightarrow N$ such that

$$\begin{aligned} p\lambda &= 1_M && \text{and} \\ \lambda p &= 1_N + d_N h + h d_N, \end{aligned}$$

which guarantee an isomorphism of homology (or cohomology), that is to say, $H_{d_M}(M) \approx H_{d_N}(N)$. SDR-data is succinctly described by

$$\left((M, d_M) \begin{array}{c} \xrightarrow{\lambda} \\ \xleftarrow{p} \end{array} (N, d_N), h \right).$$

When the inclusion map λ and the projection map p also respect additional algebraic structure on M and N , then that structure is carried over to homology (or cohomology).

Proof of Proposition 5.2. : For the deleted projective resolution M of \mathbf{B} , the non-deleted projective resolution $M^+ : M \xrightarrow{p_M} \mathbf{B} \rightarrow 0$ is acyclic, so there exists a contracting homotopy $s_i : (M^+)_i \rightarrow (M^+)_{i+1}$ where $s_{-1} : \mathbf{B} \rightarrow M_0$ such that $1_{M^+} = m_1^+ s + s m_1^+$ [Wei94]. Setting $\lambda_M = s_{-1}$ and $-(h_M)_i = s_i$ for $i \geq 0$, we produce SDR-data

$$\left((\mathbf{B}, 0) \begin{array}{c} \xrightarrow{\lambda_M} \\ \xleftarrow{p_M} \end{array} (M, m_1), h_M \right).$$

By construction, $p_M \lambda_M = 1_{\mathbf{B}}$. On M_i for $i > 0$, the map $\lambda_M p_M = 0$ and $1_{M_i} = -m_1 h_M - h_M m_1$. For $i = 0$, the identity map $1_{M_0} = -h_M m_1 + \lambda_M p_M$ and $m_1 h_M = 0$. Therefore $\lambda_M p_M = 1_M + m_1 h_M + h_M m_1$. Likewise, we can construct SDR-data

$$\left((\mathbf{sg}, 0) \begin{array}{c} \xrightarrow{\lambda_{sL}} \\ \xleftarrow{p_{sL}} \end{array} (sL, l_1), h_{sL} \right)$$

for \mathbf{sg} and sL .

We will now repeatedly “tensor” SDR-data to produce new SDR-data, using the Gugenheim tensor trick ([GLS91]) for defining the homotopy h . We illustrate this technique by showing that

$$\left((\mathbf{B} \quad \mathbf{B}, 0) \begin{array}{c} \xrightarrow{\lambda_M \quad \lambda_M} \\ \xleftarrow{p_M \quad p_M} \end{array} (M \quad M, m_1), h_M \quad M \right),$$

is SDR-data, where setting $h_M \quad M = 1_M \quad h_M + h_M \quad \lambda_M p_M$ was first proposed by Gugenheim and Lambe in [GL89]. Clearly, $(p_M \quad p_M)(\lambda_M \quad \lambda_M) = 1_{\mathbf{B} \quad \mathbf{B}}$ and it is a simple exercise to show that $1_M \quad M + m_1 h_M \quad M + h_M \quad M m_1 = (\lambda_M \quad \lambda_M)(p_M \quad p_M)$. We have verified that equation (14) and condition (iii) in definition 5.1 hold.

The tensor trick can be extended to show that

$$\left((\mathrm{T}^c(\mathbf{sg}), 0) \begin{array}{c} \xrightarrow{\mathrm{T}\lambda_{sL}} \\ \xleftarrow{\mathrm{T}p_{sL}} \end{array} (\mathrm{T}^c(sL), l_1), \mathrm{Th}_{sL} \right)$$

is SDR-data, where $\mathrm{T}\lambda_{sL} = \lambda_{sL} \quad i$ on $(\mathbf{sg}) \quad i$, where $\mathrm{T}p_{sL} = p_M \quad i$ on $(sL) \quad i$ and where $\mathrm{Th}_{sL} = \sum_{i=1}^{\infty} \mathrm{T}_i h_{sL}$ with $\mathrm{T}_i h_{sL} = \sum_{j=1}^i 1 \quad j^{-1} \quad h_{sL} \quad (\lambda_{sL} p_{sL}) \quad i-j$ on $(sL) \quad i$ (see [GLS91]). It is clear that $\mathrm{T}p_{sL} \mathrm{T}\lambda_{sL} = 1_{\mathrm{T}^c(\mathbf{sg})}$. We can show that $\mathrm{T}\lambda_{sL} \mathrm{T}p_{sL} = 1_{\mathrm{T}^c(sL)} + l_1 \mathrm{Th}_{sL} + \mathrm{Th}_{sL} l_1$ by induction on n . As a result, $\mathrm{H}_{l_1}(\mathrm{T}^c(sL)) = \mathrm{T}^c(\mathbf{sg})$. Similarly, we can define SDR-data

$$\left((\mathrm{T}^c(\mathbf{sg}) \quad \mathbf{B}, 0) \begin{array}{c} \xrightarrow{\mathrm{T}\lambda_M} \\ \xleftarrow{\mathrm{T}p_M} \end{array} (\mathrm{T}^c(sL) \quad M, m_1), \mathrm{Th}_M \right)$$

where $\mathrm{Th}_M = \sum_{i=1}^{\infty} \mathrm{T}_i h_M$ with

$$\begin{aligned} \mathrm{T}_i h_M &= \left(\sum_{j=1}^{i-1} 1 \quad j^{-1} \quad h_{sL} \quad (\lambda_{sL} p_{sL}) \quad i-j+1 \quad \lambda_M p_M \right) + 1 \quad i^{-1} \quad h_M \\ &= \mathrm{T}_{i-1} h_{sL} \quad \lambda_M p_M + 1 \quad i^{-1} \quad h_M \end{aligned}$$

on $(sL) \quad i^{-1} \quad M$.

Besides the inclusion map $i : \wedge(sL) \rightarrow \mathrm{T}^c(sL)$, there is also a splitting map $j : \mathrm{T}^c(sL) \rightarrow \wedge(sL)$ which sends $sv_{[1 \text{ to } n]}$ to $\frac{1}{n!} sv_{[1 \text{ to } n]}$. Both i and j are chain maps, so the composition $ji = 1_{\wedge(sL)}$ induces an isomorphism between $\mathrm{H}_{l_1}(\wedge(sL))$ and $\wedge(\mathrm{H}_{l_1}(sL))$, which equals $\wedge(\mathbf{sg})$. Similarly, using $(i \quad 1_M)$ and $(j \quad 1_M)$, we can show that $\mathrm{H}_{m_1}(\wedge(sL) \quad M) = \wedge(\mathbf{sg}) \quad \mathbf{B}$. Having verified that equations (12) and (13) in definition 5.1 hold, it is now straightforward to show that conditions (a) and (b) in proposition 5.2 guarantee that conditions (i) and (ii) are satisfied. \square

We can use a spectral sequence argument to prove the following result.

THEOREM 5.3. : *If the pair (M, sL) is a projective homotopy Lie-Rinehart resolution of the Lie-Rinehart pair $(\mathbf{B}, \mathfrak{sg})$, then $(\mathcal{R}, \langle D_M, \rangle)$ is a cohomological model for $(R, \langle \widehat{m}_2, \rangle)$.*

Proof of Proposition 5.3. : We have the following SDR-data:

$$\left((\bigwedge(\mathfrak{sg}), 0) \begin{array}{c} \xleftarrow{\lambda_1} \\ \xrightarrow{p_1} \end{array} (\bigwedge(sL), l_1, h_1) \right)$$

and

$$\left((\bigwedge(\mathfrak{sg}) \quad \mathbf{B}, 0) \begin{array}{c} \xleftarrow{\lambda_2} \\ \xrightarrow{p_2} \end{array} (\bigwedge(sL) \quad M, m_1), h_2 \right).$$

The E_0 term of the spectral sequence is the algebra \mathcal{R} , bigraded by the external degree of a map and the difference between the internal degree and the external degree, e.g, if the map F_n has internal degree $|F_n|$, then it has bidegree $(n, |F_n| - n)$. The differential $\langle D_M, \rangle$ splits up into the sum $\sum_{i=1}^{\infty} \langle m_i, \rangle$, each of which has bidegree $(i - 1, 2 - i)$. The map $\langle m_1, \rangle$ is the differential on E_0 . If F_n is a cocycle, then F_n is a chain map between the complexes $\bigwedge(sL)$ and $\bigwedge(sL) \quad M$. So the class of F_n induces a map $[F_n] : \bigwedge(\mathfrak{sg}) \rightarrow \bigwedge(\mathfrak{sg}) \quad \mathbf{B}$. Therefore, the E_1 term is contained in R and all elements $[F_n]$ of E_1 , have bidegree $(n, 0)$. It follows that the cohomology of \mathcal{R} is the cohomology of the E_1 term with respect to the differential $[\langle m_2, \rangle]$ given by sending $[F_n]$ to $[\langle m_2, F_n \rangle]$. The square of $[\langle m_2, \rangle]$ is zero precisely because $[\langle m_2, \langle m_2, F_n \rangle \rangle]$ equals $[-\langle m_3, \langle m_1, F_n \rangle \rangle - \langle m_1, \langle m_3, F_n \rangle \rangle]$. The first term $\langle m_3, \langle m_1, F_n \rangle \rangle$ is zero because $\langle m_1, F_n \rangle = 0$. The second term is the coboundary of $\langle m_3, F_n \rangle$, so its class with respect to $\langle m_1, \rangle$ is zero.

There is an algebra splitting from R to E_1 given by sending f_n to $[\lambda_2 f_n p_1]$, so $E_1 \approx R$ as graded commutative algebras. Furthermore, since $H_{m_1}(m_2) = \widehat{m}_2$ and $H_{l_1}(l_2) = \widehat{l}_2$, it follows that $[\langle m_2, \rangle] = \langle \widehat{m}_2, \rangle$. Therefore, E_1 and R are isomorphic as differential graded commutative algebras and have the same cohomology. \square

Constructing homotopy Lie-Rinehart resolutions for a Lie-Rinehart pair is no easy task. In an upcoming paper, we construct a homotopy Lie-Rinehart resolution for the Lie-Rinehart pair $(A/\mathcal{I}, \mathcal{I}/\mathcal{I}^2)$ which arises in the BFV formulation of classical BRST cohomology.

References

- [BF83] I. Batalin and E. Fradkin. A generalized canonical formalism and quantization of reducible gauge theories. *Physics Letters B*, 122:157–164, 1983.
- [BRS75] C. Becchi, A. Rouet, and R. Stora. Renormalization of the abelian Higgs-Kibble model. *Commun. Math. Phys.*, 42:127, 1975.
- [BV77] I. Batalin and G. Vilkovisky. Relativistic S-matrix of dynamical systems with boson and fermion constraints. *Physics Letters B*, 69:309–312, 1977.
- [BV83] I. Batalin and G. Vilkovisky. Quantization of gauge theories with linearly dependent generators. *Physics Review D*, 28:2567–2582, 1983.
- [BV85] I. Batalin and G. Vilkovisky. Existence theorem for gauge algebra. *Journal of Mathematical Physics*, 26:172–184, 1985.
- [CE48] C. Chevalley and S. Eilenberg. Cohomology theory of Lie groups and Lie algebras. *Trans. Amer. Math. Soc.*, 63:85–124, 1948.
- [dSW] A.C. do Silva, , and A. Weinstein. Lectures on geometric models for noncommutative algebras. UC-Berkeley.
- [FHST89] J. Fisch, M. Henneaux, J. Stasheff, and C. Teitelboim. Existence, uniqueness and cohomology of the classical BRST charge with ghosts of ghosts. *Commun. Math. Phys.*, 120:379, 1989.

- [FV75] E. Fradkin and G. Vilkovisky. Quantization of relativistic systems with constraints. *Physics Letters B*, 55:224–226, 1975.
- [GL89] V.K.A.M. Gugenheim and L. Lambe. Applications of perturbation theory to differential homological algebra i. *Illinois Journal of Mathematics*, 33, 1989.
- [GLS91] V.K.A.M. Gugenheim, L. Lambe, and J.D. Stasheff. Perturbation theory in differential homological algebra ii. *Illinois Journal of Mathematics*, 35(3):357–373, 1991.
- [Her72] G. C. Herz. *Cotriples and the Chevalley-Eilenberg cohomology of Lie algebra*. PhD thesis, Lehigh University, 1972.
- [HT92] M. Henneaux and C. Teitelboim. *Quantization of Gauge Systems*. Princeton University Press, 1992.
- [Kim92a] T. Kimura. Classical BRST cohomology and second class constraints. XIX International Colloquium on Group Theoretical Methods in Physics in Salamanca, Spain, July 1992.
- [Kim92b] T. Kimura. Prequantum BRST cohomology. *Contemporary Mathematics*, 132:439–457, 1992.
- [Kim93] T. Kimura. Generalized classical BRST cohomology and reduction of Poisson manifolds. *Commun. Math. Phys.*, 151:155–182, 1993.
- [KS87] B. Kostant and S. Sternberg. Symplectic reduction, BRS cohomology, and infinite-dimensional clifford algebras. *Annals of Physics*, 176:49–113, 1987.
- [Lad] T. Lada. Coderivations of cocommutative coalgebras. notes.
- [LM95] T. Lada and M. Markl. Strongly homotopy lie algebras. *Commun. in Alg.*, 23(6):2147–2161, 1995.
- [LS] T. Lada and J. D. Stasheff. Introduction to sh Lie algebras for physicists. preprint.
- [Pal61] R. S. Palais. The cohomology of Lie rings. In *Proceedings of the Symposium in Pure Mathematics, Vol. III.*, pages 130–137. American Mathematical Society, 1961.
- [Rin63] G. Rinehart. Differential forms for general commutative algebras. *Trans. Amer. Math. Soc.*, 108:195–222, 1963.
- [SS] M. Schlessinger and J. D. Stasheff. Deformation theory and rational homotopy type. to appear in Publ. Math. IHES.
- [Sta88] J. D. Stasheff. Constrained Poisson algebras and strong homotopy representations. *Bull. Amer. Math. Soc.*, pages 287–290, 1988.
- [Sta92] J. D. Stasheff. Homological (ghost) approach to constrained hamiltonian systems. *Contemp. Math.*, 132:595, 1992.
- [Sta96] J. D. Stasheff. Homological reduction of constrained Poisson algebras I: the regular case. *Journal of Differential Geometry*, ??:?, 1996.
- [Sul77] D. Sullivan. Infinitesimal computations in topology. *Publ. IHES*, 47:269–331, 1977.
- [Tat57] J. Tate. Homology of Noetherian rings and local rings. *Ill. J. Math.*, 1:14–27, 1957.
- [Tyu75] I. Tyutin. Gauge invariance in field theory and statistical physics in operator formulation (in Russian). *Lebedev Physics Inst.*, 39, 1975.
- [Wei94] C. Weibel. *An introduction to homological algebra*. Cambridge studies in advanced mathematics. Cambridge University Press, 1994.

Lars Kjeseth lkjeseth@elcamino.cc.ca.us

Mathematical Sciences Division

El Camino College

U.S.A.

Torrance

16007 Crenshaw Blvd, Torrance, CA 90506